

Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae (Coleoptera) from Turkey

Josef MERTLIK¹⁾ & Giuseppe PLATIA²⁾

1) Pohřebačka 34, CZ-533 45 Opatovice nad Labem, Czech Republic

mertlik@elateridae.com

2) Via Molino Vecchio, 21a, I-470 43 Gatteo (Forli), Italy

pinoplatia-@tele2.it

Abstract. Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey includes 477 species.

New distributional data concerning 24 species of the family Elateridae Leach, 1815 (Coleoptera) from Turkey, in accordance with Cate (2007), are provided. First records: *Adrastus samedovi* Dolin & Agajev, 1974; *Agriotes meticulus* Candèze, 1863; *Ampedus rufipennis* (Stephens 1830); *Aplotarsus angustulus* (Kiesenwetter, 1858); *Denticollis rubens* Piller et Mitterpacher, 1783; *Elathous buyssoni* Reitter, 1890; *Isidus letourneuxi* Pic, 1902; *Lacon unicolor* (Candèze, 1874); *Paracardiophorus musculus* (Erichson, 1840); *Procraterus tibialis* (Lacordaire, 1835); *Melanotus brignolii* Guglielmi & Platia, 1985; *Melanotus fulvus* Reitter, 1891; *Melanotus tenebrosus* (Erichson, 1841); *Negastrius sabulicola* (Bohemian, 1851); *Selatosomus ampliformis* Reitter, 1910; *Zorochros dermestoides* (Herbst, 1806); *Zorochros nanus* (Gurjeva, 1963) and *Zorochros quadriguttatus* (Laporte de Castelnau, 1840). Confirmed occurrence: *Agriotes lapicida* (Falderman, 1835); *Hemicrepidius niger* (Linnaeus, 1758); *Elater ferrugineus* Linnaeus, 1758; *Ischnodes sanguinicollis* (Panzer, 1793); *Plastocerus angulosus* (Germar, 1844) and *Stenagostus rosti* (Schwarz, 1897).

New distributional data concerning 2 species of the family Melasidae Leach, 1817 (Coleoptera) from Turkey, in accordance with Muona (2007), are provided. First records: *Farsus dubius* (Piller & Mitterpacher, 1783) and *Microrhagus emyi* (Rouget, 1856).

New distributional data concerning 2 species of the family Throscidae Laporte, 1840 (Coleoptera) from Turkey, in accordance with Leseigneur (2007), are provided. First records: *Trixagus atticus* Reitter, 1921 and *Trixagus duvalii* (Bonvouloir, 1859).

New synonyms: *Cardiophorus maritimus* Dolin, 1971, **syn. nov.** is synonymized with *C. vestigialis* Erichson, 1840; *Lacon aurosquamatus* (Jagemann, 1939), **syn. nov.** is synonymized with *L. lepidopterus* (Panzer, 1801); *Peripontius schimmeli* Zeising & Brunne, 2003, **syn. nov.** is synonymized with *P. crassus* (Buysson, 1906); *Selatosomus saginatus* (Ménétriés, 1832), **syn. nov.** is synonymized with *S. latus* (Fabricius, 1801).

Key words: Taxonomy, new synonym, catalogue, distribution, Coleoptera, Cebrionidae, Elateridae, Lissomidae, Melasidae, Throscidae, Turkey.

Contents

Introduction.....	2
Catalogue of the family Cebrionidae.....	2
Catalogue of the family Elateridae.....	2-27
Catalogue of the family Lissomidae.....	28
Catalogue of the family Melasidae.....	28-29
Catalogue of the family Throscidae.....	30
New distributional data of family Elateridae, Melasidae and Throscidae.....	31-36
Acknowledgements.....	36
References.....	37-40

Introduction

In this article we quote all names of species of the Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae, published from territory of Turkey. We quote 477 species from following family: Cebrionidae (1 species), Elateridae (451 species), Lissomidae (2 species), Melasidae (12 species) and Throscidae (11 species).

New distributional data concerning 24 species of the family Elateridae, 2 species of the family Melasidae and 2 species of the family Throscidae.

Abbreviations.

Citations in square brackets ([]) refer to papers recording the species from Turkey. Species preceded by an asterisk (*) indicate those of uncertain taxonomic status or with dubious Turkish records (for 34 species of the family Elateridae).

Catalogue of the family Cebrionidae Latreille, 1802

Genus *Cebrio* A. G. Olivier, 1790

Cebrio nigricollis Laporte, 1838

[Winkler 1924: 577; Lodos 1998: 1; Sánchez-Ruiz & Löbl 2007: 92].

Catalogue of the family Elateridae Leach, 1815

The classifications follows Stibick (1979) and incorporates changes made by Sánchez-Ruiz (1996).

Subfamily Agrypninae Candèze, 1857

Tribe Agrypnini Candèze, 1857

Genus *Adelocera* Latreille, 1829

Adelocera pygmaea (Baudi, 1871)

= *A. adanensis* (Jagemann, 1944)

[Jagemann 1944: 333 (as *Lacon adanensis*); Preiss & Platia 2003: 99; Cate 2007: 96].

Adelocera schuberti Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 73; Cate 2007: 96].

Genus *Agrypnus* Eschscholtz, 1829

Agrypnus crenicollis (Ménétriés, 1832)

[Bodemeyer 1900: 115 (as *Lacon*); Kesdek, Platia & Yildirim 2006: 418; Cate 2007: 97].

Agrypnus murinus (Linnaeus, 1758)

[Guglielmi & Platia 1985: 170; Lodos 1998: 46-47; Cate 2007: 97].

3 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

Agrypnus sarikamisensis Platia, Yildirim & Kesdek, 2007
[Platia, Yildirim & Kesdek, 2007].

Genus *Danosoma* C. G. Thomson, 1859

Danosoma fasciata (Linnaeus, 1758)
[Cate 2007: 100].

Genus *Lacon* Castelnau de Laporte, 1838

Lacon candezei Desbrochers des Loges, 1875

[Cate 2007: 100].

Lacon delagrangei (Buysson, 1891)

[Buysson 1891: 134; Schenkling 1925: 9 (N. Syrien, as *Adelocera*); Mertlik & Dušánek 2006: 159; Cate 2007: 100].

Lacon gillerforsi Platia & Schimmel, 1994

[Platia & Schimmel 1994: 78; Cate 2007: 101].

Lacon graecus Candèze, 1857

[Winkler 1924: 578 (as *Adelocera graeca*); Cate 2007: 101].

Lacon kapleri Platia & Schimmel, 1994

[Platia & Gudenzi 1998: 49; Cate 2007: 101].

Lacon ladae Mertlik & Dusanek, 2006

[Mertlik & Dusanek 2006: 153].

Lacon lepidopterus (Panzer, 1801)

= *L. aurosquamatus* (Jagemann, 1939) (syn. nov.) („TYPUS, *Adelocera aurosquamosa* „Jg“, Det. Jagemann; Asia Minor Amasia; Mus. Nat. Prague 26853 lav.; Lectotype *Adelocera aurosquamosa* Jagemann C.M.F. von Hayek 1974; *Lacon aurosquamosa* C.M.F. von Hayek 1974; *Lacon lepidopterus* syn. n. det. C. Wurst 1997“) (Collections of National Museum, Department of Entomology, Prague, Czech Republic).

[Jagemann 1939: 332 (as *Adelocera aurosquamosa*); Kesdek, Platia & Yildirim 2006: 419; Cate 2007: 100 (as *L. aurosquamatus*)].

Lacon punctatus (Herbst, 1779)

[Bodemeyer 1900: 115 (as *Adelocera punctata*); Schenkling 1925: 10 (as *Adelocera*); Guglielmi & Platia 1985: 170; Kesdek, Platia & Yildirim 2006: 419; Cate 2007: 101].

Lacon solai Platia & Gudenzi, 2000

[Platia & Gudenzi 2000: 66; Platia 2004: 98; Cate 2007: 102].

Lacon unicolor (Candèze, 1874)

First record for Turkey. Present paper → p. 33.

Genus *Lanelater* Arnett, 1952

Lanelater judaicus (Reiche & Saulcy, 1856)

[Guglielmi & Platia 1985: 171; Cate 2007: 102].

Lanelater persicus (Candèze, 1874)

[Platia & Gudenzi 2000: 67; Cate 2007: 102].

Tribe *Hemirhipini* Candèze, 1857

Genus *Calais* Laporte, 1838

Calais parreysi (Steven, 1830)

[Schenkling 1925: 43 (as *Alaus*); Guglielmi & Platia 1985: 171 (as *Paracalais*); Lodos 1998: 49, 50 (as *Alaus*); Kesdek, Platia & Yildirim 2006: 418; Cate 2007: 107].

Genus *Tetrigus* Candèze, 1857*Tetrigus cyprius* Baudi, 1871[Fairmaire 1893: 68 (*Ludioctenus akbesianus*); Winkler 1924: 580 (as *Ludioctenus*); Schenkling 1925: 53 (N. Syrien, as *Ludioctenus*); Preiss & Platia 2003: 103; Mertlik & Dušánek 2006: 161; Cate 2007: 108].Tribe **Oophorini** Gistel, 1856
= Conoderini Fleutiaux, 1919**Genus *Aeoloderma* Fleutiaux, 1928***Aeoloderma crucifer* (Rossi, 1790)[Bodemeyer 1900: 115 (as *Aeolus*); Guglielmi & Platia 1985: 172; Cate 2007: 103].*Aeoloderma oblitterata* (Candèze, 1859)

[Platia & Gudenzi 2000: 67; Cate 2007: 103].

Genus *Aeoloides* Schwarz, 1906* *Aeoloides atricapillus* (Germar, 1824)[Cate 2007: 105 (as *Drasterius*)].**Known from Central Asia, it is improbable that the taxon occurs in Turkey.***Aeoloides figuratus* (Germar, 1844)= *A. turcomanus* (Candèze, 1889)[Schenkling 1925: 135 (as *Drasterius*); Guglielmi & Platia 1985: 171; Cate 2007: 104].*Aeoloides griseescens* (Germar, 1844)= *A. bicarinatus* (Reitter, 1891)

[Schenkling 1925: 138; Guglielmi & Platia 1985: 172; Cate 2007: 104].

Aeoloides sarmaticus (Motschulsky, 1860)[Platia & Gudenzi 1998: 50; Cate 2007: 105 (as *Drasterius atricapillus*)].**Genus *Aeolosomus* Dolin, 1982***Aeolosomus rossii* (Germar, 1844)

[Kesdek, Platia & Yildirim 2006: 419; Cate 2007: 104].

Genus *Drasterius* Eschscholtz, 1829*Drasterius bimaculatus* (Rossi, 1790)[Bodemeyer 1900: 115; Roubal 1924: 81 (*D. bimaculatus* ab. *levantinus*); Schenkling 1925-27: 528 (as *D. bimaculatus* ab. *levantinus*); Guglielmi & Platia 1985: 171; Kesdek, Platia & Yildirim 2006: 419; Cate 2007: 105].**Subfamily Cardiophorinae Candèze, 1860**Tribe **Cardiophorini** Candèze, 1859**Genus *Cardiophorus* Eschscholtz, 1829**Subgenus *Cardiophorus* Eschscholtz, 1829*Cardiophorus acuminatus* Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 28; Cate 2007: 194].

Cardiophorus aemulus Platia & Gudenzi, 2002

5 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

- [Platia & Gudenzi 2002: 28; Cate 2007: 194].
Cardiophorus agilis Platia & Gudenzi, 2002
[Platia & Gudenzi 2002: 30; Cate 2007: 195].
* *Cardiophorus alepensis* Pic, 1914
[Lodos 1998: 49].
Known from Syria, its occurrence in Turkey is dubious.
Cardiophorus analis Schwarz, 1892
[Bodemeyer 1900: 116; Winkler 1924: 592; Schenkling 1925: 220; Lodos 1998: 49; Platia & Gudenzi 2002: 47; Kesdek, Platia & Yildirim 2006: 419; Cate 2007: 195].
Cardiophorus anticus Erichson, 1840
[Platia & Gudenzi 2000: 601, Kabalak & Sert 2005: 56; Cate 2007: 195].
Cardiophorus apterus Platia & Gudenzi, 2002
[Platia & Gudenzi 2002: 31; Cate 2007: 195].
Cardiophorus araxicola Iablokoff-Khnzorian, 1970
[Platia 2004: 107; Cate 2007: 195].
Cardiophorus bellus Platia & Gudenzi, 2000
[Platia & Gudenzi 2000: 603; Cate 2007: 195].
Cardiophorus carnosus Platia & Gudenzi, 2002
[Platia & Gudenzi 2002: 32; Cate 2007: 196].
Cardiophorus cyanipennis Mulsant & Wachanru, 1852
[Winkler 1924: 590; Schenkling 1925: 224; Guglielmi & Platia 1985: 214; Lodos 1998: 49; Platia & Gudenzi 2000: 601, Kabalak & Sert 2005: 56; Cate 2007: 196].
Cardiophorus discicollis (Herbst, 1806)
[Bodemeyer 1900: 116; Horion 1953: 227; Lodos 1998: 48, 49; Platia & Gudenzi 2000: 601, Kabalak & Sert 2005: 56; Kesdek, Platia & Yildirim 2006: 419; Cate 2007: 196].
Cardiophorus dolini Mardjanian, 1985
[Platia & Gudenzi 2002: 48; Cate 2007: 196].
Cardiophorus ebeninus (Germar, 1824)
[Cate 2007: 196].
Cardiophorus eliasi Pic, 1904
[Platia & Gudenzi 2000: 602; Cate 2007: 196].
Cardiophorus fossulatus Platia & Gudenzi, 2002
[Platia & Gudenzi 2002: 32; Cate 2007: 197].
Cardiophorus frequens Platia & Gudenzi, 2002
[Platia & Gudenzi 2002: 34; Cate 2007: 197].
Cardiophorus gramineus (Scopoli, 1763)
[Platia & Gudenzi 2000: 602; Cate 2007: 197].
Cardiophorus hayekae Platia & Gudenzi, 2002
[Platia & Gudenzi 2002: 35; Cate 2007: 197].
Cardiophorus hinkei Frivaldszky, 1837
[Platia & Gudenzi 2000: 602; Preiss & Platia 2003: 104; Cate 2007: 197].
Cardiophorus imitator Platia & Gudenzi, 2002
[Platia & Gudenzi 2002: 36; Cate 2007: 197].
Cardiophorus impressiventris Schwarz, 1900
= *C. boyadjiani* Pic, 1908
[Winkler 1924: 593 (as *C. boyadjianii* and *C. impressiventris*); Schenkling 1925: 223 (as *C. boyadjianii*); Schenkling 1925: 226; Guglielmi & Platia 1985: 214; Lodos 1998: 49 (as *C. boyadjianii*); Platia & Gudenzi 2002: 48; Cate 2007: 197].
* *Cardiophorus insignis* Desbrochers, 1875
[Lodos 1998: 49].
Known from Israel, Syria and „Palestine“, its occurrence in Turkey is dubious.
Cardiophorus kindermannii Candèze, 1860
= *C. akbesianus* Pic, 1899
[Bodemeyer 1900: 116; Winkler 1924: 590; Schenkling 1925: 220 (as *C. akbesianus*); Lodos 1998: 49; Platia & Gudenzi 2000: 602; Cate 2007: 198].
Cardiophorus kinzelbachi Chassain, 1979

[Platia & Gudenzi 2002: 48; Cate 2007: 198].

Cardiophorus levis Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 36; Cate 2007: 198].

Cardiophorus lundbergi Platia & Gudenzi, 2000

[Platia & Gudenzi 2000: 604; Cate 2007: 198].

Cardiophorus maculicollis Reiche & Saulcy, 1856

[Platia & Gudenzi 2000: 602; Cate 2007: 198].

* *Cardiophorus marmottani* Bulysson, 1912

[Schenkling 1925: 237 (as *Perrinellus*); Lodos 1998: 49].

Known from North Africa (Morocco), it is improbable that the taxon occurs in Turkey.

Cardiophorus megathorax (Faldermann, 1835)

= *C. antennalis* Desbrochers, 1875

[Guglielmi & Platia 1985: 214 (as *C. antennalis*); Lodos 1998: 49 (as *C. antennalis*); Platia & Gudenzi 2002: 49; Kesdek, Platia & Yildirim 2006: 420; Cate 2007: 198].

Cardiophorus miniaticollis Candèze, 1860

[Platia & Gudenzi 2000: 602; Cate 2007: 198].

Cardiophorus neuberti Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 38; Cate 2007: 199].

Cardiophorus nigerrimus Erichson, 1840

[Lodos 1998: 49; Platia & Gudenzi 2002: 49; Cate 2007: 199].

Cardiophorus nigratissimus Bulysson, 1891

[Bulysson 1891: 134; Guglielmi & Platia 1985: 214; Lodos 1998: 49; Platia & Gudenzi 2002: 49, Kabalak & Sert 2005: 57; Kesdek, Platia & Yildirim 2006: 420; Cate 2007: 199].

Cardiophorus parvulus Platia & Gudenzi, 2000

[Platia & Gudenzi 2000: 142; Platia & Gudenzi 2002: 49; Kesdek, Platia & Yildirim 2006: 420; Cate 2007: 199].

Cardiophorus pavesii Platia & Gudenzi, 2000

[Platia & Gudenzi 2000: 604; Cate 2007: 199].

Cardiophorus picinus Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 39; Cate 2007: 199].

* *Cardiophorus picticollis* Kraatz, 1882

[Winkler 1924: 590; Schenkling 1925: 228; Lodos 1998: 49].

Known from Central Asia (Tajikistan, Uzbekistan), it is improbable that the taxon occurs in Turkey.

Cardiophorus plurimus Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 39; Cate 2007: 199].

Cardiophorus procerus Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 40; Cate 2007: 199].

Cardiophorus proprius Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 42; Cate 2007: 199].

Cardiophorus ressli Platia & Gudenzi, 2000

[Platia & Gudenzi 2000: 605; Cate 2007: 200].

Cardiophorus rotundicollis Frivaldszky, 1845

= *C. turgescens* Bulysson, 1891

[Schenkling 1925: 232 (as *C. turgescens*); Lodos 1998: 49 (as *C. turgescens*); Platia & Gudenzi 2002: 50; Cate 2007: 200].

* *Cardiophorus ruficollis* (Linnaeus, 1758)

[Lodos 1998: 49].

Euro-Siberian species, its occurrence in Turkey is dubious.

Cardiophorus ruficruris (Brullé, 1832)

= *C. tibialis* Erichson, 1840

[Platia & Gudenzi 2002: 50; Preiss & Platia, 2003: 104; Cate 2007: 200].

* *Cardiophorus rufipes* Goeze, 1777

= *C. goezei* Sánchez-Ruiz, 1996

[Schenkling 1925: 230; Cate 2007: 200].

Known from western Mediterranean, it is improbable that the taxon occurs in Turkey.

Cardiophorus sacratus Erichson, 1840

7 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

[Schenkling 1925: 231 (*C. sacratus* ab. *tokatensis* Pic, 1913); Guglielmi & Platia 1985: 214; Lodos 1998: 49; Platia & Gudenzi 2000: 603; Preiss & Platia 2003: 104, Kabalak & Sert 2005: 57; Kesdek, Platia & Yildirim 2006: 420; Cate 2007: 200].

Cardiophorus samai Platia & Gudenzi, 2000

[Platia & Gudenzi 2000: 605; Cate 2007: 200].

Cardiophorus schuberti Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 42; Cate 2007: 200].

Cardiophorus schurmanni Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 44; Cate 2007: 200].

Cardiophorus somcheticus Schwarz, 1896

[Cate 2007: 200].

Cardiophorus stussineri Bulyssson, 1913

[Platia & Gudenzi 2002: 51; Preiss & Platia 2003: 104; Cate 2007: 200].

Cardiophorus syriacus (Linnaeus, 1758)

[Winkler 1924: 590; Schenkling 1925: 231; Guglielmi & Platia 1985: 214; Lodos 1998: 49; Platia & Gudenzi 2000: 603; Kesdek, Platia & Yildirim 2006: 420; Cate 2007: 200].

* *Cardiophorus tenellus* Reiche, 1856

[Bodemeyer 1900: 116; Schenkling 1925: 232, Lodos 1998: 49].

Known from Egypt, Israel, Jordan, Lebanon and Syria, its occurrence in Turkey is dubious.

Cardiophorus turgidus Erichson, 1840

[Lodos 1998: 49; Platia & Gudenzi 2002: 51; Cate 2007: 201].

Cardiophorus uncinatus Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 46; Cate 2007: 201].

Cardiophorus varius Cate, Platia & Schimmel, 2002

[Cate, Platia & Schimmel 2002: 51].

Cardiophorus vestigialis Erichson, 1840

= *C. atramentarius* Erichson, 1840

= *C. erichsoni* Bulyssson, 1901

= *C. maritimus* Dolin, 1971 (syn. nov.)

= *C. nijazi* Dolin & Agajew, 1983

[Bodemeyer 1900: 116 (as *C. rufipes*); Horion 1953: 230 (as *C. erichsoni*); Guglielmi & Platia 1985: 215; Lodos 1998: 49 (as *C. atramentarius*, *C. erichsoni* and as *C. rufipes*); Platia & Gudenzi 2002: 51; Preiss & Platia 2003: 104; Kabalak & Sert 2005: 56 (as *C. maritimus*); Kesdek, Platia & Yildirim 2006: 420; Cate 2007: 198 (as *C. maritimus*); Cate 2007: 201].

Cardiophorus winkleri Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 47].

Cardiophorus witzgalli Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 46; Cate 2007: 201].

Cardiophorus zeisingi Platia, 2004

[Platia 2004: 106; Cate 2007: 201].

Subgenus *Perrinellus* Bulyssson, 1899

* *Cardiophorus aegyptiacus* Candèze, 1860

[Farmaire 1866; Lodos 1998: 49].

Known from Egypt and Tunisia, it is improbable that the taxon occurs in Turkey.

Genus *Dicronychus* Brullè, 1832

Dicronychus adanensis (Pic, 1908)

[Winkler 1924: 594 (as *Platynychus*); Schenkling 1925: 233 (as *Platynychus*); Lodos 1998: 49 (as *Cardiophorus*); Platia & Gudenzi 2004: 18; Cate 2007: 203].

Dicronychus apteriformis Platia & Gudenzi, 2004

[Platia & Gudenzi 2004: 10; Cate 2007: 203].

Dicronychus bicolor Platia & Gudenzi, 2004

[Platia & Gudenzi 2004: 10; Cate 2007: 204].

- Dicronychus bozdaghensis* Platia & Gudenzi, 2004
 [Platia & Gudenzi 2004: 10; Cate 2007: 204].
- Dicronychus cinereus* (Herbst, 1784)
 [Guglielmi & Platia 1985: 212; Platia & Gudenzi 2004: 18; Kabalak & Sert 2005: 57; Cate 2007: 204].
- Dicronychus decorus* (Faldermann, 1835)
 [Bodemeyer 1900: 116 (as *Cardiophorus*); Winkler 1924: 594 (as *Platynychus*); Schenkling 1925: 234 (as *Platynychus*); Lodos 1998: 49 (as *Cardiophorus*); Platia & Gudenzi 2004: 18; Cate 2007: 204].
- Dicronychus deminutus* Platia & Gudenzi, 2004
 [Platia & Gudenzi 2004: 12; Cate 2007: 204].
- Dicronychus desbrochersi* Platia & Gudenzi, 2004
 [Platia & Gudenzi 2004: 12; Cate 2007: 204].
- * *Dicronychus drurei* (Pic, 1908)
 [Lodos 1998: 49].
- Known from Iraq, its occurrence in Turkey is dubious.**
- Dicronychus equiseti* (Herbst, 1784)
 [Winkler 1924: 594 (as *Platynychus*); Schenkling 1925: 234 (as *Platynychus*); Horion 1953: 237; Lodos 1998: 49; Platia & Gudenzi 2004: 18; Cate 2007: 204].
- * *Dicronychus extinctus* (Erichson, 1840)
 [Lodos 1998: 49 (as *Cardiophorus*)].
- Known from Egypt and „Mesopotamia“, its occurrence in Turkey is dubious.**
- Dicronychus freudei* Platia & Gudenzi, 2007
 [Platia & Gudenzi 2007: 93].
- Dicronychus gillerforsii* Platia & Gudenzi, 2004
 [Platia & Gudenzi 2004: 12; Cate 2007: 204].
- Dicronychus iconiensis* (Pic, 1908)
 [Winkler 1924: 594 (as *Platynychus*); Schenkling 1925: 235 (as *Platynychus*); Guglielmi & Platia 1985: 212 (as *D. incanus*); Lodos 1998: 49 (as *Cardiophorus*); Platia & Gudenzi 2004: 18; Cate 2007: 204].
- Dicronychus merkli* (Pic, 1910)
 [Winkler 1924: 595 (as *Platynychus*); Schenkling 1925: 235 (as *Platynychus*); Guglielmi & Platia 1985: 212; Lodos 1998: 49; Preiss & Platia 2003: 104; Platia & Gudenzi 2004: 20; Cate 2007: 205].
- Dicronychus mutilus* Platia & Gudenzi, 2004
 [Platia & Gudenzi 2004: 14; Cate 2007: 205].
- Dicronychus obscuripennis* (Pic, 1899)
 [Winkler 1924: 594 (as *Platynychus decorus* ab. *obscuripennis*); Schenkling 1925: 234 (as *Platynychus decorus* ab. *obscuripennis*); Platia & Gudenzi 2004: 20; Cate 2007: 205].
- Dicronychus pici* Platia & Gudenzi, 2004
 [Platia & Gudenzi 2004: 14; Cate 2007: 205].
- Dicronychus rubripes* (Germar, 1824)
 [Platia & Gudenzi 2004: 20; Kabalak & Sert 2005: 57; Kesdek, Platia & Yildirim 2006: 420; Cate 2007: 205].
- Dicronychus senaci* Desbrochers des Loges, 1870
 [Winkler 1924: 595 (as *Platynychus*); Platia & Martini 1990: 46; Platia & Gudenzi 2004: 21; Cate 2007: 205].
- Dicronychus turcicus* Platia & Gudenzi, 2004
 [Platia & Gudenzi 2004: 16; Cate 2007: 205].
- Dicronychus variatus* Desbrochers, 1875
 [Guglielmi & Platia 1985: 214; Platia & Gudenzi 2004: 21; Cate 2007: 205].
- * *Dicronychus versicolor* (Mulsant & Guillebeau, 1856)
 [Bodemeyer 1900: 116 (as *Cardiophorus*); Lodos 1998: 49; Cate 2007: 205].
- Known from South Europe (France, Italy, Slovenia, Greece), its occurrence in Turkey is dubious.**

Genus *Paracardiophorus* Schwarz, 1895

- Paracardiophorus musculus* (Erichson, 1840)
First records for Turkey. Present paper → p. 33.

9 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

Genus *Ryukyucardiophorus* Ôhira, 1973

Ryukyucardiophorus mertliki Platia & Gudenzi, 1999
[Platia & Gudenzi 1999: 23; Cate 2007: 207].

Subfamily Dendrometrinae Gistel, 1856

= Denticollinae Stein & J. Weise, 1877

Tribe Dendrometrini Gistel, 1856

Subtribe **Athouina** Candèze, 1859

Genus *Athous* Eschscholtz, 1829

Subgenus *Athous* Eschscholtz, 1829

Athous haemorrhoidalis (Fabricius, 1801)

[Bodemeyer 1900: 116; Horion 1953: 258; Yüksel 1970: 35; Lodos 1998: 43, 44; Preiss & Platia 2003: 103; Kabalak & Sert 2005: 51; Kesdek, Platia & Yıldırım 2006: 421; Cate 2007: 156].

Athous nadoraz Mertlik & Dusanek 2006

[Mertlik & Dušánek 2006: 146; Dušánek & Mertlik 2007: 69, 72].

Athous vittatus (Fabricius, 1792)

= *A. vittatus* (Gmelin, 1790)

[Reitter 1905: 46; Winkler 1924: 601; Schenkling 1925-1927: 323 (*Athous vittatus* ab. *denigratus*); Horion 1953: 258; Guglielmi & Platia 1985: 173; Lodos 1998: 43; Kesdek, Platia & Yıldırım 2006: 421; Cate 2007: 157].

Subgenus *Euplathous* Reitter, 1905

* *Athous proximus* (Hampe, 1864)

[Reitter 1905: 80; Winkler 1924: 603; Schenkling 1925-1927: 334; Lodos 1998: 43].

Known from Bulgaria, Greece and Romania, its occurrence in Turkey is dubious.

Subgenus *Haplatheous* Reitter, 1905

* *Athous carpathicus* Reitter, 1889

[Kabalak & Sert 2005: 51].

Known from Carpathian Mountains (Romania), it is improbable that the taxon occurs in Turkey.

Athous marginicollis Reitter, 1890

[Guglielmi & Platia 1985: 177; Lodos 1998: 43; Cate 2007: 159].

Athous subfuscus (O.F. Müller, 1764)

[Guglielmi & Platia 1985: 179; Lodos 1998: 43; Kabalak & Sert 2005: 51; Kesdek, Platia & Yıldırım 2006: 421; Cate 2007: 159].

Athous turcicus Reitter, 1905

[Reitter 1905: 75; Winkler 1924: 603; Schenkling 1925-1927: 330; Lodos 1998: 43; Cate 2007: 159].

Subgenus *Orthathous* Reitter, 1905

Athous acutangulus Fairmaire, 1866

= *A. thessalicus* Reiche, 1869

[Reitter 1905: 88; Winkler 1924: 603; Schenkling 1925-1927: 334, 340 (as *A. thessalicus*); Lodos 1998: 43; Platia & Gudenzi 1996: 240; Cate 2007: 160].

Athous anatolicus Platia, 1989

- [Platia 1989: 82; Platia & Gudenzi 1996: 240; Cate 2007: 160].
* *Athous apfelbecki* Reitter, 1905
[Lodos 1998: 43].
Known from Bulgaria, Hungary, Romania, Serbia and Montenegro, its occurrence in Turkey is dubious.
- Athous artvinensis* Platia, Yildirim & Kesdek, 2007
[Platia, Yildirim & Kesdek, 2007].
- Athous audisioi* Guglielmi & Platia, 1985
[Guglielmi & Platia 1985: 180; Lodos 1998: 43; Platia & Gudenzi 1996: 240; Cate 2007: 160].
- Athous barriesi* Platia & Gudenzi, 1996
[Platia & Gudenzi 1996: 236; Cate 2007: 160].
- Athous benedicti* Platia, 2003
[Platia 2003: 263; Cate 2007: 160].
- * *Athous bicolor* (Goeze, 1777)
[Lodos 1998: 43 (as *A. longicornis*)].
European species reaching eastward to Ukraine, it is improbable that the taxon occurs in Turkey.
- Athous cavazzutii* Platia, 2003
[Platia 2003: 264; Cate 2007: 161].
- Athous daccordii* Guglielmi & Platia, 1985
[Guglielmi & Platia 1985: 182; Lodos 1998: 43; Platia & Gudenzi 1996: 240; Cate 2007: 161].
- Athous dasycerus* Bulysson, 1890
[Reitter 1905: 101; Winkler 1924: 604; Schenckling 1925-1927: 338; Guglielmi & Platia 1985: 183; Lodos 1998: 43; Platia & Gudenzi 1996: 240; Cate 2007: 161].
- Athous demirsoyi* Platia, Kabalak & Sert, 2007
[Platia, Kabalak & Sert 2007: 45].
- Athous dusaneki* Platia, 2003
[Platia 2003: 265; Dušánek & Mertlik 2007: 69, 72; Cate 2007: 161].
- Athous edirnensis* Platia & Gudenzi, 2000
[Platia & Gudenzi 2000: 71; Cate 2007: 161].
- Athous epiurus* Stierlin, 1875
[Cate 2007: 161].
- Athous fragariae* Platia & Kovancı, 2005
[Platia & Kovancı 2005: 5].
- Athous freudei* Platia, 1989
[Platia & Gudenzi 2000: 74; Cate 2007: 161].
- Athous frontalis* Platia & Schimmel, 1991
[Platia & Schimmel 1991: 136; Platia & Gudenzi 1996: 240; Cate 2007: 161].
- Athous gianassoi* Platia & Gudenzi, 1996
[Platia & Gudenzi 1996: 236; Cate 2007: 161].
- Athous graecus* Platia, 1989
[Platia & Gudenzi 2000: 74; Cate 2007: 161].
- Athous gudenzii* Guglielmi & Platia, 1985
[Guglielmi & Platia 1985: 183; Platia & Gudenzi 1996: 240; Lodos 1998: 43; Cate 2007: 161].
- Athous hajekii* Platia & Gudenzi, 2005
[Platia & Gudenzi 2005: 120].
- Athous hetzeli* Platia, 2004
[Platia 2004: 99; Cate 2007: 161].
- Athous kovancii* Platia, 2003
[Platia 2003: 265; Cate 2007: 162].
- Athous lassallei* Platia & Gudenzi, 1996
[Platia & Gudenzi 1996: 237; Platia 2003: 266; Cate 2007: 162].
- Athous malmusii* Platia & Gudenzi, 2000
[Platia & Gudenzi 2000: 72; Cate 2007: 162].
- Athous margheritae* Guglielmi & Platia, 1985
[Guglielmi & Platia 1985: 184; Platia & Gudenzi 1996: 240; Lodos 1998: 43; Dušánek & Mertlik 2007: 68, 72; Cate 2007: 162].

11 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

- Athous mertliki* Platia & Gudenzi, 2002
[Platia & Gudenzi 2002: 83; Cate 2007: 162].
- Athous paflagonensis* Platia & Gudenzi, 1998
[Platia & Gudenzi 1998: 632; Cate 2007: 162].
- Athous ponticus* Platia & Gudenzi, 2007
[Platia & Gudenzi 2007: 84].
- Athous propinquus* Buysson, 1889
[Reitter 1905: 101; Lodos 1998: 43; Cate 2007: 162].
- Athous ruffoi* Guglielmi & Platia, 1985
[Guglielmi & Platia 1985: 186; Platia & Gudenzi 1996: 240; Lodos 1998: 43; Cate 2007: 162].
- Athous sabatinellii* Guglielmi & Platia, 1985
[Guglielmi & Platia 1985: 187; Platia & Gudenzi 1996: 240; Lodos 1998: 43; Cate 2007: 162].
- Athous sameki* Platia, 2003
[Platia 2003: 266; Cate 2007: 162].
- Athous senaci* Buysson, 1889
= *A. acutangulus* var. *ottomanus* Reitter, 1905
[Winkler 1924: 603 (as *A. acutangulus* var. *ottomanus*), 604; Schenkling 1925-1927: 340; Platia & Martini 1990: 34; Platia & Gudenzi 1996: 240; Lodos 1998: 43; Cate 2007: 162].
- Athous snizeki* Platia, 2004
[Platia 2004: 100; Cate 2007: 162].
- Athous svihlai* Platia & Gudenzi, 1998
[Platia & Gudenzi 1998: 632; Cate 2007: 162].
- Athous tekkirazicus* Platia, 2003
[Platia 2003: 266; Dušánek & Mertlik 2007: 69, 72; Cate 2007: 162].
- Athous triberti* Guglielmi & Platia, 1985
[Guglielmi & Platia 1985: 189; Platia & Gudenzi 1996: 240; Lodos 1998: 43; Cate 2007: 163].
- Athous warchałowskii* Platia & Tarnawski, 1998
[Platia & Tarnawski 1998: 270; Cate 2007: 163].
- Athous wewalkai* Platia, 1989
[Platia 1989: 81; Platia & Gudenzi 1996: 240; Cate 2007: 163].
- Athous zanettii* Guglielmi & Platia, 1985
[Guglielmi & Platia 1985: 190; Platia & Gudenzi 1996: 240; Lodos 1998: 43; Cate 2007: 163].
- Athous zбузеки* Platia & Gudenzi, 2007
[Platia & Gudenzi 2007: 83].

Genus *Hemicrepidius* Germar, 1839

- Hemicrepidius corvinus* (Reitter, 1905)
[Reitter 1905: 38; Guglielmi & Platia 1985: 193; Cate 2007: 170].
- Hemicrepidius hirtus* (Herbst, 1784)
[Schenkling 1925-1927: 313; Horion 1953: 255; Yüksel 1970: 36 (as *Athous*); Guglielmi & Platia 1985: 193; Preiss & Platia 2003: 104; Kesdek, Platia & Yıldırım 2006: 421; Cate 2007: 170].
- * *Hemicrepidius koenigi* Schwarz, 1897
[Schenkling 1925-1927: 314 (Kurdistan); Lodos 1998: 43 (as *Athous*)].
- Known from Armenia, its occurrence in Turkey is dubious.**
- Hemicrepidius niger* (Linnaeus, 1758)
[Kabalak & Sert 2005: 52].
- Confirmed occurrence in Turkey. Present paper → p. 31.**
- Hemicrepidius nigritulus* (Reitter, 1890)
[Guglielmi & Platia 1985: 194; Kesdek, Platia & Yıldırım 2006: 421; Cate 2007: 170].
- Hemicrepidius schneideri* (Kiesenwetter, 1877)
[Guglielmi & Platia 1985: 194; Kesdek, Platia & Yıldırım 2006: 421; Cate 2007: 171].
- Hemicrepidius vulpeculus* (Reitter, 1890)
[Guglielmi & Platia 1985: 194; Kesdek, Platia & Yıldırım 2006: 421; Cate 2007: 171].

Genus *Pseudocrepidophorus* Dolin et Agajev, 1988

Pseudocrepidophorus concavifrons Wurst, 1994

[Wurst, 1994: 59; Cate 2007: 172].

Pseudocrepidophorus flavesiensis (Eschscholtz, 1818)

[Guglielmi & Platia 1985: 173; Dušánek & Mertlik 2007: 69, 72; Cate 2007: 172].

Genus *Stenagostus* C. G. Thomson, 1859

Stenagostus rhombeus (Olivier, 1790)

[Horion 1953: 253; Mertlik & Dušánek 2006: 160; Cate 2007: 172].

Stenagostus rosti (Schwarz, 1897)

[Orlov 1989: ?; Lodos 1998: 43].

Confirmed occurrence in Turkey. Present paper → p. 34.

Stenagostus zuercheri Reitter, 1909

[Winkler 1924: 599; Schenkling 1925-1927: 313; Platia & Gudenzi 1996: 142; Lodos 1998: 43; Mertlik & Dušánek 2006: 161; Kesdek, Platia & Yildirim 2006: 421; Cate 2007: 172].

Subtribe **Dendrometrina** Gistel, 1856

Genus *Cidnopus* C. G. Thomson, 1859

Cidnopus crassipes (Schwarz, 1900)

[Winkler 1924: 598 (Syr. b.); Winkler 1924: 597 (as *Limonius*); Schwarz 1900: 105; Reitter 1905: 15; Schenkling 1925-1927: 292; Cate & Platia; 1989: 187; Cate 2007: 164].

Cidnopus pilosus (Leske, 1785)

[Schenkling 1925-1927: 294; Horion 1953: 244; Guglielmi & Platia 1985: 172; Cate & Platia 1989: 187; Cate 2007: 164].

Cidnopus schurmanni Platia & Gudenzi, 1998

[Mertlik & Dušánek 2006: 158; Cate 2007: 164].

Cidnopus turcicus Platia, 2004

[Platia 2004: 40; Cate 2007: 164].

Genus *Elathous* Reitter, 1890

Elathous buyssoni Reitter, 1890

First record for Turkey. Present paper → p. 32.

Elathous depressus Platia & Schimmel, 1991

[Platia & Schimmel 1991: 131; Cate 2007: 164].

Elathous lizlerwernerii Mertlik & Dusanek, 2006

[Mertlik & Dušánek 2006: 152].

Elathous mertlici Platia & Schimmel, 1992

[Platia & Schimmel 1992: 93; Cate 2007: 164].

Elathous nigricans Platia & Schimmel, 1991

[Platia & Schimmel 1991: 132; Kesdek, Platia & Yildirim 2006: 421; Cate 2007: 164].

Elathous pavesii Platia & Schimmel, 1992

[Platia & Schimmel 1992: 94; Cate 2007: 164].

Elathous rufobasalis Wurst, 1994

[Wurst, 1994: 62; Cate 2007: 164].

Elathous schwarzi Reitter, 1905

[Reitter 1905: 22; Winkler 1924: 598 (as *Leptoschema*); Schenkling 1925-1927: 303; Cate 2007: 164].

Elathous smyrnensis Reitter, 1905

[Reitter 1905: 23; Winkler 1924: 598 (as *Leptoschema*); Schenkling 1925-1927: 303; Cate 2007: 164].

Elathous turcicus Platia & Gudenzi, 2005

[Platia & Gudenzi 2005: 114].

13 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

Genus *Limoniscus* Reitter, 1905

Limoniscus elegans (Buysson, 1891)

[Reitter 1905: 15; Schenkling 1925-1927: 297 (N. Syrien); Platia & Gudenzi 1996: 141; Chassain, 1998: 33; Cate 2007: 165].

Limoniscus violaceus (Ph.Müller, 1843)

[Mertlik & Dušánek 2006: 159; Jansson & Coskun 2006; Cate 2007: 165]

Genus *Limonius* Eschscholtz, 1829

* *Limonius araratensis* Faldermann, 1835

[Faldermann 1835: 168; Winkler 1924: 598; Schenkling 1925-1927: 292].

The taxonomic status of this taxon is unresolved (Cate 2007: 209, as nomina dubia).

Limonius minutus (Linnaeus, 1758)

[Winkler 1924: 597 (as *Limonius*); Schenkling 1925-1927: 292; Horion 1953: 245; Guglielmi & Platia 1985: 172; Kabalak & Sert 2005: 52; Cate 2007: 165].

Limonius poneli Leseigneur et Mertlik, 2007

[Leseigneur & Mertlik 2007: 232].

Genus *Nothodes* LeConte, 1861

Nothodes parvulus (Panzer, 1799)

[Bodemeyer 1900: 116 (as *Limonius*); Winkler 1924: 597 (as *Limonius*); Horion 1953: 246; Guglielmi & Platia 1985: 172; Cate 2007: 165].

Genus *Solskyana* Dolin, 1978

* *Solskyana villiger* (Solsky, 1881)

[Winkler 1924: 597 (as *Limonius*)].

Known from Armenia, Kyrgyzstan, Kazakhstan and Uzbekistan, its occurrence in Turkey is dubious.

Subtribe **Denticollina** Stein et J. Wiese, 1877

Genus *Denticollis* Piller et Mitterpacher, 1783

Denticollis linearis (Linnaeus, 1758)

[Winkler 1924: 615; Schenkling 1925-27: 496; Horion 1953: 308; Cate 2007: 167].

Denticollis parallellicollis Aubé, 1850

[Guglielmi & Platia 1985: 193; Cate 2007: 168].

Denticollis rubens Piller et Mitterpacher, 1783

First record for Turkey. Present paper → p. 32.

Tribe **Prosternini** Gistel, 1856

= *Ctenicerini* Fleutiaux, 1936

Genus *Anostirus* C. G. Thomson, 1859

Anostirus alexandri Wurst, 1995

[Wurst 1995: 152; Cate 2007: 174].

Anostirus ataturki Platia & Gudenzi, 2000

[Platia & Gudenzi 2000: 68; Kesdek, Platia & Yildirim 2006: 421; Cate 2007: 174].

Anostirus eschscholtzi (Faldermann, 1835)

[Gurjeva 1989: 141; Kesdek, Platia & Yildirim 2006: 422; Cate 2007: 174].

Anostirus gracilicollis (Stierlin, 1896)

- [Kesdek, Platia & Yildirim 2006: 422; Cate 2007: 174].
Anostirus holtzi (Schwarz, 1902)
[Schwarz 1902: 289; Winkler 1924: 605; Schenkling 1925-1927: 366; Cate 2007: 174].
Anostirus incostatus (Pic, 1905)
[Platia & Gudenzi 1998: 50; Cate 2007: 174].
Anostirus lederi (Heyden, 1878)
[Bodemeyer 1900: 116 (as *Ludius*); Gurjeva 1989: 137, Tarnawski 1996: 618; Cate 2007: 174].
Anostirus nubilosus Wurst & Schimmel, 1995
[Wurst & Schimmel 1995: 25; Cate 2007: 174].
Anostirus purpureus (Poda, 1761)
[Cate 2007: 175].
Anostirus saltinii Cate, Platia & Schimmel, 2002
[Schenkling 1925-1927: 367 (?as *A. sulphuripennis*); Mertlik & Dušánek 2006: 158; Cate 2007: 175].
Anostirus trivialis Gurjeva, 1988
[Mertlik & Dušánek 2006: 158; Cate 2007: 175].
Anostirus turcicus Platia & Mertlik, 1996
[Platia & Mertlik 1996: 225; Cate 2007: 175].

Genus *Aplotarsus* Stephens, 1830

- Aplotarsus angustulus* (Kiesenwetter, 1858)
First record for Turkey. Present paper → p. 31.

Genus *Ctenicera* Latreille, 1829

- Ctenicera pectinicornis* (Linnaeus, 1758)
[Guglielmi & Platia 1985: 194; Kesdek, Platia & Yildirim 2006: 422; Cate 2007: 177].

Genus *Hypoganus* Kiesenwetter, 1863

- Hypoganus inunctus* (Panzer, 1795)
= *H. cinctus* (Paykull, 1800)
[Pic 1903: 129 (as *H. cinctus* ab. *akbesianus*); Schenkling 1925-1927: 403 (as *H. cinctus* ab. *akbesianus*); Cate 2007: 177].
Hypoganus stepanovi Denisova, 1948
[Platia & Gudenzi 1999: 26; Cate 2007: 177].

Genus *Neopristilophus* Buysson, 1894

- * *Neopristilophus insitivus* (Germar, 1824)
= *N. depressus* Germar, 1822
[Schenkling 1925-1927: 384; Horion 1953: 285; Lodos 1998: 40 (as *Selatosomus*)].
Submediterranean species, its occurrence in Turkey is dubious.

Genus *Prosternon* Latreille, 1834

- Prosternon syriacum* (Buysson, 1891)
= *P. anatolicum* Reitter, 1901
[Bodemeyer 1900: 116 (as *Ludius*); Winkler 1924: 609 (as *P. anatolicum* and *P. syriacum*); Schenkling 1925-1927: 399 (as *P. anatolicum* and *P. syriacum*); Guglielmi & Platia 1985: 195; Lodos 1998: 40 (as *Selatosomus*); Kesdek, Platia & Yildirim 2006: 422; Cate 2007: 180].
Prosternon tessellatum (Linnaeus, 1758)
[Guglielmi & Platia 1985: 194; Kesdek, Platia & Yildirim 2006: 422; Cate 2007: 180].

15 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

Genus *Pseudanostirus* Dolin, 1964

Pseudanostirus jasoni Dolin & Chantladze, 1987
[Platia & Gudenzi 1996: 140; Cate 2007: 180].

Genus *Selatosomus* Stephens, 1830

Subgenus *Pristilophus* Latreille, 1834

Selatosomus melancholicus alpestris (Fabricius, 1798)
[Gurjeva 1989: 231; Mertlik 2000: 8; Kesdek, Platia & Yildirim 2006: 422; Cate 2007: 181].

Subgenus *Selatosomus* Stephens, 1830

Selatosomus aeneus (Linnaeus, 1758)
[Guglielmi & Platia 1985: 195; Lodos 1998: 40, 42; Cate 2007: 181].

Selatosomus amplicollis (Germar, 1843)
[Guglielmi & Platia 1985: 195; Lodos 1998: 40; Mertlik 2000: 8; Kesdek, Platia & Yildirim 2006: 422; Cate 2007: 182].

Selatosomus armeniacus Dolin, 1982
[Gurjeva 1989: 258; Cate 2007: 182].

* *Selatosomus atratus* (Ballion, 1878)
[Schenkling 1925-1927: 374; Lodos 1998: 40].

Known from Kyrgyzstan and Kazakhstan, it is improbable that the taxon occurs in Turkey.

Selatosomus beysehiricus Mertlik, 2000
[Guglielmi & Platia 1985: 195 (as *S. latissimus*); Mertlik 2000: 4; Cate 2007: 182].

Selatosomus caucasicus (Ménétriés, 1879)
[Bodemeyer 1900: 116 (as *Ludius violaceus*); Lodos 1998: 40 (as *Selatosomus violaceus*); Mertlik 2000: 9;
Cate 2007: 182].

Selatosomus latissimus Reitter, 1910
[Reitter 1910: 169; Winkler 1924: 607; Schenkling 1925-1927: 376; Gurjeva 1989: 256; Guglielmi & Platia
1985: 195 (as *S. latissimus* - nec Reitter 1910); Lodos 1998: 40; Mertlik 2000: 10; Cate 2007: 182].

Selatosomus latus (Fabricius, 1801)
= *S. gravidus* (Germar, 1843)
= *S. saginatus* (Ménétriés, 1832) (**syn. nov.**)
[Bodemeyer 1900: 116; Schenkling 1925-1927: 374 (as *S. corpulentus*); Guglielmi & Platia 1985: 196;
Gurjeva 1989: 256 (as *S. saginatus*); Lodos 1998: 40, 41; Mertlik 2000: 10, Kabalak & Sert 2005: 52; Kesdek,
Platia & Yildirim 2006: 422 (as *S. gravidus*); Cate 2007: 182 (as *S. latus* and *S. saginatus*)].

Selatosomus pecirkanus Reitter, 1910
[Mertlik 2000: 9; Cate 2007: 182].

Selatosomus punctipennis Reitter, 1910
[Reitter 1910: 169; Winkler 1924: 607; Schenkling 1925-1927: 374; Gurjeva 1989: 247; Lodos 1998: 40;
Mertlik 2000: 9; Cate 2007: 182].

Selatosomus sameki Mertlik, 2000
[Guglielmi & Platia 1985: 195 (as *S. latissimus*); Mertlik 2000: 6; Cate 2007: 182].

Subgenus *Warchałowskianus* Tarnawski, 1995

Selatosomus ampliformis Reitter, 1910
First record for Turkey. Present paper → p. 34.

Selatosomus barlaensis Mertlik, 2000
[Mertlik 2000: 2; Cate 2007: 183].

Subfamily Elaterinae Leach, 1815

Tribe Agriotini Champion, 1894

Genus *Agriotes* Eschscholtz, 1829

Subgenus *Agriotes* Eschscholtz, 1829

Agriotes acuminatus (Stephens, 1830)

[Schenkling 1925-1927: 443 (as *Agriodrastus*); Horion 1953: 295; Guglielmi & Platia 1985: 202; Lodos 1998: 16; Cate 2007: 114].

Agriotes adanensis Pic, 1910

[Winkler 1924: 611 (as *Ectinus*); Schenkling 1925-1927: 441 (as *Ectinus*); Gurjeva 1972: 868; Lodos 1998: 16; Platia 2003: 24; Cate 2007: 114].

Agriotes adlbaueri Platia & Schimmel, 1992

[Platia & Schimmel 1992: 91; Platia & Gudenzi 1998: 55; Cate 2007: 114].

Agriotes anatolicus Platia, 2003

[Platia 2003: 14; Cate 2007: 114].

Agriotes aquilus Platia, 2003

[Platia 2003: 15; Cate 2007: 114].

Agriotes audisioi Platia & Schimmel, 1992

[Platia & Schimmel 1992: 90; Cate 2007: 114].

Agriotes barriesi Cate & Platia, 1997

[Cate & Platia 1997: 109; Platia 2003: 24; Cate 2007: 114].

Agriotes brevis Candèze, 1863

[Schenkling 1925-1927: 445; Horion 1953: 299; Gurjeva 1972: 872; Lodos 1998: 16, 17, 35; Cate 2007: 114].

Agriotes cilicensis Platia & Gudenzi, 1998

[Platia & Gudenzi 1998: 634; Cate 2007: 114].

Agriotes colonnelli Guglielmi & Platia, 1985

[Guglielmi & Platia 1985: 202; Lodos 1998: 16; Platia & Gudenzi 1998: 56; Cate 2007: 114].

Agriotes conspicuus Schwarz, 1891

[Winkler 1924: 611 (as *Ectinus*); Schenkling 1925-1927: 442 (as *Ectinus*); Gurjeva 1972: 868; Lodos 1998: 16; Platia 2003: 23; Cate 2007: 114].

Agriotes constrictus Reitter, 1900

[Winkler 1924: 611 (as *Ectinus*); Schenkling 1925-1927: 442 (as *Ectinus*); Gurjeva 1972: 868; Lodos 1998: 16; Cate 2007: 114].

Agriotes crassiusculus Platia & Gudenzi, 1998

[Platia & Gudenzi 1998: 634; Cate 2007: 114].

Agriotes cribrithorax Pic, 1910

[Winkler 1924: 611 (as *Ectinus*); Schenkling 1925-1927: 442 (as *Ectinus*); Gurjeva 1972: 866; Platia & Schimmel 1994: 82; Lodos 1998: 16; Cate 2007: 114].

Agriotes defreinai Platia & Gudenzi, 1998

[Platia & Gudenzi 1998: 57; Platia 2003: 24; Cate 2007: 114].

Agriotes depressus Schwarz, 1891

[Platia & Gudenzi 1998: 634; Cate 2007: 114].

Agriotes dusaneki Platia & Gudenzi, 1998

[Platia & Gudenzi 1998: 633; Platia 2003: 24; Cate 2007: 114].

Agriotes furlani Platia, 2003

[Platia 2003: 16; Cate 2007: 115].

* *Agriotes ganglbaueri* Schwarz, 1891

[Lodos 1998: 16].

Known from Lebanon and Syria, its occurrence in Turkey is dubious.

Agriotes goksunensis Platia, 2004

[Platia 2004: 101; Cate 2007: 115].

Agriotes graecus Franz, 1967

17 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

- [Platia & Gudenzi 1999: 27; Cate 2007: 115].
Agriotes granulosus Platia & Schimmel, 1992
[Platia & Schimmel 1992: 90; Platia & Gudenzi 1998: 636; Cate 2007: 115].
Agriotes gurgistanus (Falderman, 1835)
[Schenkling 1925-1927: 446; Gurjeva 1972: 867; Guglielmi & Platia 1985: 204; Lodos 1998: 16, 17, 35; Platia & Gudenzi 1998: 634; Kesdek, Platia & Yildirim 2006: 422; Cate 2007: 115].
Agriotes heydeni Schwarz, 1891
[Winkler 1924: 610 (as *Ectinus*); Schenkling 1925-1927: 443 (as *Ectinus*); Gurjeva 1972: 868; Lodos 1998: 16; Platia 2003: 23; Cate 2007: 115].
Agriotes informis Schwarz, 1891
[Winkler 1924: 611; Schenkling 1925-1927: 446; Lodos 1998: 16; Platia 2003: 23; Cate 2007: 115].
Agriotes infuscatus Desbrochers des Loges, 1870
[Gurjeva 1972: 868; Guglielmi & Platia 1985: 204; Lodos 1998: 16, 17; Cate 2007: 115].
Agriotes integracollis Reitter, 1911
[Schenkling 1925-1927: 443 (as *A. acuminatus* ab. *integracollis*); Cate 2007: 115].
Agriotes izmirensis Cate & Platia, 1997
[Cate & Platia 1997: 111; Platia 2003: 24; Cate 2007: 115].
Agriotes karsantianus Pic, 1910
[Gurjeva 1972: 869 (*A. reitteri* Schwarz, 1891); Platia & Schimmel 1993: 54; Cate 2007: 115].
Agriotes kraatzii Schwarz, 1891
[Winkler 1924: 610 (as *Ectinus*); Schenkling 1925-1927: 443 (as *Ectinus*); Gurjeva 1972: 868; Platia & Gudenzi 1997: 209; Lodos 1998: 16, 17; Platia 2003: 24; Cate 2007: 115].
Agriotes lapicida (Falderman, 1835)
First record for Turkey. Present paper → 30.
Agriotes leinfestii Platia & Gudenzi, 1998
[Platia & Gudenzi 1998: 57; Platia 2003: 24; Kesdek, Platia & Yildirim 2006: 422; Cate 2007: 115].
Agriotes lineatus (Linnaeus, 1767)
[Bodemeyer 1900: 116; Schenkling 1925-1927: 447; Horion 1953: 300; Yüksel 1970: 28; Gurjeva 1972: 871; Guglielmi & Platia 1985: 204; Lodos 1998: 11, 16, 17, 19, 20, 31, 32; Kesdek, Platia & Yildirim 2006: 422; Cate 2007: 115].
* *Agriotes litigiosus* (Rossi, 1792)
[Lodos 1998: 16, 35].
Known from Austria, Bulgaria, France (Corse), Greece, Italy, Macedonia, Slovenia, Spain and Switzerland, its occurrence in Turkey is dubious.
Agriotes lizleri Platia, 2003
[Platia 2003: 17; Cate 2007: 116].
Agriotes lundbergi Platia, 1989
[Platia 1989: 83; Preiss & Platia 2003: 106; Cate 2007: 116].
Agriotes mertlici Platia, 2003
[Platia 2003: 17; Cate 2007: 116].
Agriotes meticulosus Candèze, 1863
[Gurjeva 1972: 872; Lodos 1998: 16, 35 (as *A. meticulosus* Cand.)].
Confirmed occurrence in Turkey. Present paper → p. 30.
* *Agriotes modestus* Kiesenwetter, 1858
= *A. incognitus* Schwarz, 1891
= *A. ponticus* Stepanov, 1935
[Schenkling 1925-1927: 450; Lodos 1998: 16].
Submediterranean species, its occurrence in Turkey is dubious.
Agriotes nigropubens Reitter, 1904
[Winkler 1924: 611; Schenkling 1925-1927: 451; Gurjeva 1972: 873; Lodos 1998: 16; Platia & Gudenzi 1998: 636; Kesdek, Platia & Yildirim 2006: 422; Cate 2007: 116].
Agriotes nigror Platia, 2003
[Platia 2003: 20; Cate 2007: 116].
Agriotes nuceus Fairmaire, 1866
[Winkler 1924: 611 (as *Ectinus*); Schenkling 1925-1927: 443 (as *Ectinus*); Gurjeva 1972: 868; Guglielmi & Platia 1985: 204; Lodos 1998: 16; Platia & Gudenzi 1998: 56; Platia 2003: 22; Cate 2007: 116].

Agriotes obscurus (Linnaeus, 1758)

[Yüksel 1970: 30; Lodos 1998: 11, 16, 17, 25, 27, 30; Kesdek, Platia & Yildirim 2006: 422].

Agriotes paludum Kiesenwetter, 1859

[Bodemeyer 1900: 116; Guglielmi & Platia 1985: 204; Lodos 1998: 16; Kesdek, Platia & Yildirim 2006: 422; Cate 2007: 116].

Agriotes petterssoni Platia & Gudenzi, 2000

[Platia & Gudenzi 2000: 74; Cate 2007: 116].

Agriotes pilosellus (Schönherr, 1817)

[Yüksel 1970: 24 (*A. pilosus*); Lodos 1998: 16, Lodos 1998: 16 (*A. pilosus*); Kesdek, Platia & Yildirim 2006: 423; Cate 2007: 116].

Agriotes propleuralis Platia & Gudenzi, 1998

[Platia & Gudenzi 1998: 635; Cate 2007: 117].

Agriotes proximus Schwarz, 1891

= *A. lineatoides* Reitter, 1911

[Reitter 1911: 222; Schenckling 1925-1927: 450 (as *A. lineatus* ab. *proximus*); Guglielmi & Platia 1985: 205, Lodos 1998: 16; Kabalak & Sert 2005: 54; Kesdek, Platia & Yildirim 2006: 423; Cate 2007: 117].

* *Agriotes radula* Desbrochers des Loges, 1875

= *Agriotes brevicollis* Schwarz, 1891

[Cate 2007: 117].

Known from Lebanon, Syria and Israel, its occurrence in Turkey is dubious.

* *Agriotes reitteri* Schwarz, 1891

[Winkler 1924: 612; Lodos 1998: 16].

Known from Rossia (South European Territory) and Georgia, its occurrence in Turkey is dubious.

Agriotes riesei Platia, 2004

[Platia, 2004: 42; Cate 2007: 117].

Agriotes rufipalpis (Brullé, 1832)

= *A. rubiginosus* Candèze, 1863

[Guglielmi & Platia 1985: 205; Lodos 1998: 16; Lodos 1998: 16 (as *A. rubiginosus*); Kesdek, Platia & Yildirim 2006: 423].

Agriotes sameki Platia, 2003

[Platia 2003: 21; Cate 2007: 117].

Agriotes schuberti Platia, 2004

[Platia 2004: 104; Cate 2007: 117].

Agriotes schurmanni Platia & Gudenzi, 1998

[Platia & Gudenzi 1998: 58; Platia 2003: 25; Cate 2007: 117].

Agriotes sexualis Platia, 2004

[Platia 2004: 104; Cate 2007: 117].

Agriotes solai Platia, 2004

[Platia 2004: 42; Cate 2007: 117].

* *Agriotes sordidus* (Illiger, 1807)

[Lodos 1998: 16, 17 & 18; Kabalak & Sert 2005: 54].

Known from western Mediterranean, it is improbable that the taxon occurs in Turkey.

Agriotes sputator (Linnaeus, 1758)

[Bodemeyer 1900: 116; Schenckling 1925-1927: 455; Horion 1953: 299; Yüksel 1970: 27, 43; Gurjeva 1972: 872; Guglielmi & Platia 1985: 205; Lodos 1998: 11, 16-20, 25, 34, 41; Kesdek, Platia & Yildirim 2006: 423; Cate 2007: 117].

Agriotes starcki Schwarz, 1891

[Schenckling 1925-1927: 457; Lodos 1998: 16; Platia & Gudenzi 1999: 27; Cate 2007: 118].

Agriotes strigosus Kiesenwetter, 1859

[Platia & Gudenzi 1999: 28; Preiss & Platia 2003: 106; Cate 2007: 118].

Agriotes subsulcatus Pic, 1913

[Winkler 1924: 611 (as *Ectinus*); Schenckling 1925-1927: 443 (as *Ectinus*); Gurjeva 1972: 868; Lodos 1998: 16; Platia 2003: 24; Cate 2007: 118].

Agriotes sylviae Cate & Platia, 1997

[Cate & Platia, 1997: 111; Cate 2007: 118].

Agriotes tauricus Heyden, 1882

19 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

[Guglielmi & Platia 1985: 205; Lodos 1998: 16; Cate 2007: 118].

Agriotes tristiculus Reitter, 1906

[Winkler 1924: 612; Schenkling 1925-1927: 457; Gurjeva 1972: 874; Guglielmi & Platia 1985: 205; Lodos 1998: 16, 33, 34; Kesdek, Platia & Yildirim 2006: 423; Cate 2007: 118].

Agriotes turcicus Candèze, 1863

[Bodemeyer 1900: 116; Schenkling 1925-1927: 457; Gurjeva 1972: 869; Cate 2007: 118].

Agriotes ustulatus (Schaller, 1783)

[Guglielmi & Platia 1985: 205; Lodos 1998: 16; Kesdek, Platia & Yildirim 2006: 423; Cate 2007: 118].

Agriotes wernerii Platia, 2003

[Platia 2003: 22; Cate 2007: 118].

Agriotes witzgalli Platia & Schimmel, 1993

[Platia & Schimmel 1993: 52; Cate 2007: 118].

Genus *Agriotes* species incertae sedis

* *Agriotes aterrimus* Heyden

[Lodos 1998: 16].

Probably mistaken transcription of correct name.

* *Agriotes lapicriola* Falderman

[Lodos 1998: 35].

Probably mistaken transcription of correct name.

Genus *Dalopius* Eschscholtz, 1829

Dalopius marginatus (Linnaeus, 1758)

[Guglielmi & Platia 1985: 206; Cate 2007: 119].

Dalopius radiculosus Gurjeva, 1964

[Mertlik & Dušánek 2006: 159; Cate 2007: 119].

Genus *Ectinus* Eschscholtz, 1829

Ectinus aterrimus (Linnaeus, 1761)

[Lodos 1998: 16 (as *Agriotes aterrimus* Payk.)].

Genus *Pseudagriotes* Schwarz, 1896

Pseudagriotes holtzi Schwarz, 1896

[Winkler 1924: 613; Schenkling 1925-1927: 464; Platia & Gudenzi 1996: 148; Cate 2007: 120].

Tribe Ampedini Gistel, 1856

Genus *Ampedus* Dejean, 1833

Ampedus adlbaueri Schimmel, 1990

[Schimmel 1990: 76; Cate 2007: 120].

Ampedus aethiops (Lacordaire, 1835)

= *A. scrofa* Germar, 1844

[Cate 2007: 121].

Ampedus agajewi Dolin, 1970

[Platia & Gudenzi 1999: 29; Cate 2007: 121].

Ampedus agilis Platia, 2001

[Platia 2001: 66; Cate 2007: 121].

Ampedus anatolicus Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 86; Cate 2007: 121].

Ampedus balteatus (Linnaeus, 1785)

- [Cate 2007: 121].
- Ampedus canaliculatus* (Reitter, 1918)
[Reitter 1918: 86; Winkler 1924: 582 (as *Elater*); Schenkling 1925: 142 (as *Elater*); Cate 2007: 122].
- Ampedus cinnaberinus* (Eschscholtz, 1829)
[Reitter 1918: 85 (*Elater cinnaberinus* sbsp. *atropilosus*); Schenkling 1925: 143 (as *Elater*); Guglielmi & Platia 1985: 196; Kesdek, Platia & Yildirim 2006: 423; Cate 2007: 122].
- Ampedus circassicus* (Reitter, 1887)
[Reitter 1918: 98; Platia & Gudenzi 1999: 29; Kesdek, Platia & Yildirim 2006: 424; Cate 2007: 122].
- Ampedus decumanus* Gurjeva, 1977
[Platia & Gudenzi 1999: 30; Kesdek, Platia & Yildirim 2006: 424; Cate 2007: 122].
- Ampedus demaisonii* (Buysson, 1898)
[Winkler 1924: 583 (as *Elater*); Schenkling 1925: 144 (as *Elater*); Guglielmi & Platia 1985: 196; Cate 2007: 122].
- Ampedus elegantulus* (Schönherr, 1817)
[Winkler 1924: 584 (as *Elater*); Schenkling 1925: 144 (as *Elater*); Horion 1953: 190; Kabalak & Sert 2005: 53; Kesdek, Platia & Yildirim 2006: 424; Cate 2007: 122].
- Ampedus elongatulus* (Fabricius, 1787)
= *A. glycereus* (Herbst, 1784)
[Winkler 1924: 583 (as *Elater elongatulus* ab. *bodemeyeri*); Schenkling 1925: 145 (as *Elater elongatulus* ab. *bodemeyeri*); Platia & Gudenzi 2000: 74 (as *A. glycereus*); Cate 2007: 123].
- Ampedus erythrogonus* (P. W. Müller, 1821)
[Cate 2007: 123].
- Ampedus ganglbaueri* (Reitter, 1889)
[Reitter 1918: 96; Platia & Gudenzi 1999: 30; Cate 2007: 123].
- Ampedus hirticollis* (Semenov, 1891)
[Bodemeyer 1900: 115; Cate 2007: 123].
- Ampedus lubricus* (Buysson 1891)
[Buysson 1891: 135; Winkler 1924: 583 (Syr. b., as *Elater*); Schenkling 1925: 146 (N. Syrien, as *Elater*); Cate 2007: 124].
- Ampedus nigerrimus* (Lacordaire, 1835)
[Cate 2007: 125].
- Ampedus nigrinus* (Herbst, 1784)
[Cate 2007: 125].
- Ampedus nigroflavus* (Goeze, 1777)
[Horion 1953: 187; Cate 2007: 125].
- Ampedus ocellatus* (Buysson, 1891)
= *A. fragarinus* Wurst, 1995
[Buysson 1891: 135; Reitter 1918: 88; Winkler 1924: 582 (Syr. b., as *Elater*); Schenkling 1925: 148 (N. Syrien, as *Elater*); Wurst 1995: 149; Wurst 1997: 1; Cate 2007: 125].
- Ampedus ochropterus* Germar, 1844
[Horion 1953: 184; Kesdek, Platia & Yildirim 2006: 424; Cate 2007: 125].
- Ampedus platiai* Schimmel, 1990
[Schimmel 1990: 78; Cate 2007: 126].
- Ampedus pomonae* (Stephens, 1830)
[Cate 2007: 126].
- Ampedus pomorum* (Herbst in Füssly, 1784)
= *Ampedus robustus* Bouwer, 1980
[Guglielmi & Platia 1985: 196 (as *Ampedus robustus*); Kabalak & Sert 2005: 53; Cate 2007: 126].
- Ampedus praeustus* (Fabricius, 1792)
[Cate 2007: 126].
- Ampedus pulcher* (Baudi, 1871)
[Reitter 1918: 97; Platia & Gudenzi 1999: 30, Preiss & Platia, 2003: 106; Cate 2007: 126].
- Ampedus punctatus* Schwarz, 1893
[Reitter 1918: 86; Winkler 1924: 582 (as *Elater*); Schenkling 1925: 152 (as *Elater*); Cate 2007: 126].
- Ampedus ressli* Platia & Gudenzi, 1999
[Platia & Gudenzi 1999: 29; Cate 2007: 127].

21 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

Ampedus rufipennis (Stephens, 1830)

First records for Turkey. Present paper → p. 31.

Ampedus samedovi Dolin & Ajajew, 1983

[Platia & Gudenzi 1999: 30; Cate 2007: 127].

Ampedus sanguineus (Linnaeus, 1758)

[Bodemeyer 1900: 115; Horion 1953: 184; Cate 2007: 127].

Ampedus sanguinolentus (Schrank, 1776)

[Laibner 2000; Kabalak & Sert 2005: 53; Cate 2007: 127].

Ampedus schoettlei Zeising & Brunne, 2003

[Zeising & Brunne 2003: 192; Cate 2007: 127].

Ampedus sinuatus Germar, 1844

[Bodemeyer 1900: 115; Reitter 1918: 99; Horion 1953: 195; Platia & Gudenzi 1999: 118; Kesdek, Platia & Yildirim 2006: 424; Cate 2007: 127].

Ampedus turcicus Platia, Kabalak & Sert, 2007

[Platia, Kabalak & Sert 2007: 44].

Ampedus wachtangi Dolin, 1970

[Guglielmi & Platia 1985: 196 (det. as *A. balteatus*); Platia & Gudenzi 1999: 30; Kesdek, Platia & Yildirim 2006: 424; Cate 2007: 128].

Genus *Brachygonus* Buysson, 1912

Brachygonus frater Wurst, 1995

[Wurst 1995: 147; Cate 2007: 129].

Brachygonus gratiosus Platia & Schimmel, 1991

[Platia & Schimmel 1991: 139; Cate 2007: 129].

Brachygonus hadullanus Wurst, 1995

[Wurst 1995: 31; Cate 2007: 129].

Brachygonus megerlei (Lacordaire, 1835)

[Mertlik & Dušánek 2006: 158; Cate 2007: 129].

Brachygonus ruficepsoides Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 87; Cate 2007: 129].

Genus *Haterumelater* Ôhira, 1968

Haterumelater fulvago (Marseul, 1868)

= *H. julianum* Penev, 1987

= *H. preissi* Schimmel, 1995

= *H. picinus* (Buysson, 1891)

[Winkler 1924: 585 (as *Ischnodes*); Schenkling 1925: 165 (as *Ischnodes picinus*); Guglielmi & Platia 1985: 198 (as *Ischnodes picinus*); Preiss & Platia 2003: 107; Cate 2007: 130].

Haterumelater languidus (Buysson, 1891)

[Winkler 1924: 585 (as *Ischnodes*); Schenkling 1925: 165 (as *Ischnodes*); Preiss & Platia; 2003: 110; Cate 2007: 130].

Genus *Ischnodes* Germar, 1844

Ischnodes kalleci Mertlik, 2005

[Mertlik 2005: 167; Cate 2007: 130].

Ischnodes sanguinicollis (Panzer, 1793)

[Platia 1994: 281].

Confirmed occurrence in Turkey. Present paper → p. 32.

Genus *Reitterelater* Platia et Cate, 1990

Reitterelater dubius Platia & Cate, 1990

[Platia & Cate 1990: 113; Cate 2007: 130].

Tribe **Elaterini** Leach, 1815

Genus *Elater* Linnaeus, 1758

Elater ferrugineus ferrugineus Linnaeus, 1758

[Yüksel 1970: 31 (as *Ludius*)].

Confirmed occurrence in Turkey. Present paper → p. 32.

Elater tauricus (Schwarz, 1897)

[Bodemeyer 1900: 116 (as *Melanotus*); Winkler 1924: 610 (as *Ludius*); Schenkling 1925-1927: 428 (as *Ludius*); Wurst 1994: 118; Cate 2007: 132].

Genus *Mulsanteus* Gozis, 1875

= *Neotrichophorus* Jacobson, 1913

Mulsanteus guillebeaui (Mulsant & Godart, 1853)

= *M. guillebelli* (Mulsant & Godart, 1853)

[Winkler 1924: 610 (as *Neotrichophorus*); Schenkling 1925-1927: 432 (as *Neotrichophorus*); Guglielmi & Platia 1985: 206 (as *Neotrichophorus*); Preiss & Platia 2003: 104; Kesdek, Platia & Yildirim 2006: 424; Cate 2007: 133 (as *N. guillebeaui*)].

Mulsanteus manuelae Platia & Gudenzi, 1998

[Platia & Gudenzi 1998: 59; Cate 2007: 133].

Mulsanteus schaumi (Candèze, 1881)

[Winkler 1924: 610 (as *Neotrichophorus*); Schenkling 1925-1927: 432 (as *Neotrichophorus*); Guglielmi & Platia 1985: 206 (as *Neotrichophorus*); Kesdek, Platia & Yildirim 2006: 424; Cate 2007: 133].

Genus *Pittonotus* Jasquelin du Val, 1860

* *Pittonotus simoni* (Stierlin, 1879)

[Winkler 1924: 610; Schenkling 1925-1927: 431 (as *P. theseus* var. *simoni*)].

Known from Syria, its occurrence in Turkey is dubious.

Pittonotus theseus (Germar, 1817)

[Bodemeyer 1900: 116 (as *Ectinus*); Winkler 1924: 610; Guglielmi & Platia 1985: 206; Lodos 1998: 36 (as *Melanotus thseus* Germ.); Preiss & Platia 2003: 104; Kesdek, Platia & Yildirim 2006: 424; Cate 2007: 133].

Tribe **Megapenthini** Gurjeva, 1973

Genus *Ectamenogonus* Buysson, 1894

Ectamenogonus montandoni (Buysson, 1888)

[Platia 1994: 283; Kubáň 1995: 25; Preiss & Platia, 2003: 112; Mertlik & Dušánek 2006: 159; Cate 2007: 136].

Genus *Megapenthes* Kiesenwetter, 1858

Megapenthes lugens (Redtenbacher, 1842)

[Mertlik & Dušánek 2006: 160; Cate 2007: 137].

Megapenthes rutilipennis Candèze, 1859

[Cate 2007: 137].

Genus *Procraerus* Reitter, 1905

Procraerus bicolor Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 89; Cate 2007: 138].

Procraerus carinifrons (Desbrochers des Loges, 1875)

[Kesdek, Platia & Yildirim 2006: 425].

23 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

Procraerus levantinus Platia & Gudenzi, 2007

[Platia & Gudenzi, 2007: 88].

Procraerus tibialis (Lacordaire, 1835)

First record for Turkey. Present paper → p. 34.

Tribe **Physorhinini** Candèze, 1859

Genus *Podeonius* Kiesenwetter, 1858

Podeonius acuticornis (Germar, 1824)

[Mertlik & Dušánek 2006: 160; Cate 2007: 140].

Genus *Porthmidius* Germar, 1847

Porthmidius austriacus (Schrank, 1781)

[Winkler 1924: 586; Horion 1953: 209; Cate 2007: 140].

Tribe **Pomachiliini** Candèze, 1859

Genus *Betarmon* Kiesenwetter, 1858

Betarmon bisbimaculatus (Fabricius, 1803)

[Horion 1953: 211; Platia 1994: 306; Cate 2007: 140].

Genus *Idolus* Desbrochers, 1875

Idolus adrastoides (Reitter, 1888)

[Guglielmi & Platia 1985: 202; Cate 2007: 140].

Genus *Idotarmonides* Agajev, 1985

Idotarmonides anatolicus (Candèze, 1881)

[Winkler 1924: 587 (as *Idolus*); Schenkling 1925: 196 (as *Idolus*); Cate 2007: 141].

Idotarmonides rydhi Platia & Schimmel, 1992

[Platia & Schimmel 1992: 87; Cate 2007: 141].

Genus *Isidus* Mulsant et Rey, 1874

Isidus letourneuxi Pic, 1902

First record for Turkey. Present paper → p. 33.

Isidus moreli Mulsant & Rey, 1874

[Winkler 1924: 598; Preiss & Platia 2003: 104; Cate 2007: 141].

Tribe **Synaptini** Gistel, 1856
= Adrastini Candèze, 1863

Genus *Adrastus* Eschscholtz, 1829

Adrastus anatolicus Platia & Schimmel, 1991

[Platia & Schimmel 1991: 138; Kesdek, Platia & Yildirim 2006: 425; Cate 2007: 108].

Adrastus circassicus Reitter, 1896

[Guglielmi & Platia 1985: 198; Cate 2007: 108].

Adrastus dolini Welschmied, 1978

[Platia & Gudenzi 1998: 55; Cate 2007: 108].

Adrastus limbatus (Fabricius, 1776)

[Schenkling 1925-1927: 491; Horion 1953: 303; Guglielmi & Platia 1985: 200; Cate 2007: 108].

Adrastus longicornis Gurjeva, 1976

[Platia & Gudenzi 1998: 55; Cate 2007: 108].

Adrastus montanus (Scopoli, 1763)

[Yüksel 1970: 47; Guglielmi & Platia 1985: 200; Platia & Gudenzi 1998: 55; Cate 2007: 108].

Adrastus pallens (Fabricius, 1792)

[Schenkling 1925-1927: 493; Schenkling 1925-27; Horion 1953: 304; Cate 2007: 108].

Adrastus protractus Roubal, 1924

[Guglielmi & Platia 1985: 200; Cate 2007: 109].

Adrastus rachifer (Fourcroy, 1785)

[Yüksel 1970: 49; Guglielmi & Platia 1985: 200; Cate 2007: 109].

Adrastus samedovi Dolin & Agajev, 1974

First records for Turkey. Present paper → p. 30.

Genus *Peripontius* Gurjeva, 1979

Peripontius concolor (Desbrochers des Loges, 1875)

[Winkler 1924: 614 (as *Silesis concolor* a. *juvencus* Buysson, 1887); Schenkling 1925-1927: 488 (as *Silesis concolor* ab. *juvencus*); Cate 2007: 111].

Peripontius crassus (Buysson, 1906)

= *P. schimmeli* Zeising & Brunne, 2003: 193 (**syn. nov.**)

[Platia 2003: 90; Zeising & Brunne 2003: 193 (as *P. schimmeli*); Cate 2007: 111 (as *P. crassus* and *P. schimmeli*)].

Peripontius dentatus Platia & Schimmel, 1991

[Bodemeyer 1900: 116 (as *P. concolor* and *P. juvencus*); Guglielmi & Platia 1985 (as *P. concolor*); Platia & Schimmel 1991: 137; Cate 2007: 111].

Peripontius elegantulus Platia, 2003

[Platia 2003: 85; Cate 2007: 111].

Peripontius jagemannii Platia & Gudenzi, 2000

[Cate 2007: 111].

Peripontius oertzeni Schwarz, 1897

[Cate 2007: 111].

Peripontius omissus (Buysson, 1889)

[Platia & Martini 1990; Platia & Schimmel 1991: 138; Kesdek, Platia & Yildirim 2006: 425; Cate 2007: 111].

Peripontius saroveci Platia, 2003

[Platia 2003: 89; Cate 2007: 111].

Peripontius terminatus (Erichson, 1842)

[Winkler 1924: 614; Horion 1953: 303; Guglielmi & Platia 1985: 202; Preiss & Platia 2003: 114, Kabalak & Sert 2005: 54; Cate 2007: 111].

Peripontius turcicus Platia & Tarnawski, 1998

[Platia & Tarnawski 1998: 271; Cate 2007: 111].

Genus *Synaptus* Eschscholtz, 1829

Synaptus filiformis (Fabricius, 1781)

[Bodemeyer 1900: 116; Winkler 1924: 614; Schenkling 1925-1927: 483; Horion 1953: 302; Yüksel 1970: 45; Guglielmi & Platia 1985: 198; Kesdek, Platia & Yildirim 2006: 426; Cate 2007: 113].

Genus *Tolphorea* Gurjeva, 1983

Tolphorea kurdistanicus Platia & Gudenzi, 1996

[Platia & Gudenzi 1996: 142; Cate 2007: 113].

Tolphorea pallida Platia, 2004

[Platia 2004: 105; Cate 2007: 113].

25 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

Tolphorea turcica Platia & Gudenzi, 2000
[Platia & Gudenzi 2000: 76; Cate 2007: 113].

Subfamily Hypnoidinae Schwarz, 1906

Genus *Hypnoidus* Dillwyn, 1829

Hypnoidus anatolicus Platia & Gudenzi, 2000
[Platia & Gudenzi 2000: 67; Cate 2007: 154].

Subfamily Melanotinae Candèze, 1859

Genus *Melanotus* Eschscholtz, 1829

Subgenus *Melanotus* Eschscholtz, 1829

Melanotus aemulus (Erichson, 1841)
[Schwarz 1892: 153; Schenkling 1925-1927: 272; Guglielmi & Platia 1985: 206; Lodos 1998: 36; Cate 2007: 141].

Melanotus armeniacus Schwarz, 1891
[Lodos 1998: 36].

Melanotus atricapillus Reitter, 1891
[Cate 2007: 142].

Melanotus bajulus (Erichson, 1841)
[Kabalak & Sert 2006: 216; Kesdek, Platia & Yildirim 2006: 426; Cate 2007: 142].

Melanotus brignolii Guglielmi & Platia, 1985

First records for Turkey. Present paper → p. 33.

Melanotus brunneipes (Germar, 1824)
[Schenkling 1925-1927: 272; Guglielmi & Platia 1985: 209; Lodos 1998: 35, 36, 38, 39; Kesdek, Platia & Yildirim 2006: 426; Cate 2007: 142].

Melanotus castanipes (Paykull, 1800)
[Schenkling 1925-27: 273; Lodos 1998: 36; Cate 2007: 142].

Melanotus crassicollis (Erichson, 1841)
[Horion 1953: 241; Platia 1994: 317; Lodos 1998: 36; Cate 2007: 143].

Melanotus dichrous (Erichson, 1841)
[Schwarz 1892: 162; Schenkling 1925-1927: 275; Guglielmi & Platia 1985: 209; Lodos 1998: 36; Cate 2007: 143].

Melanotus fraseri Platia & Schimmel, 1993
[Platia & Schimmel 1993: 55; Cate 2007: 144].

Melanotus fulvus Reitter, 1891

First records for Turkey. Present paper → p. 33.

Melanotus fusciceps (Gyllenhal, 1817)
= *M. robustus* (Erichson, 1841)
[Schwarz 1892: 163; Schenkling 1925-1927: 276; Guglielmi & Platia 1985: 209; Guglielmi & Platia 1985: 211 (as *M. robustus*); Platia & Martini 1990: 44; Lodos 1998: 35, 36, 39; Lodos 1998: 36 (as *M. robustus*); Preiss & Platia, 2003: 104; Kesdek, Platia & Yildirim 2006: 426; Cate 2007: 144].

Melanotus monticola (Ménétriés, 1832)
[Guglielmi & Platia 1985: 209; Lodos 1998: 36; Kesdek, Platia & Yildirim 2006: 427; Cate 2007: 147].

* *Melanotus mutatus* Harold, 1869
[Lodos 1998: 36].

Known from Iran and Syria, its occurrence in Turkey is dubious.

Melanotus osellai Guglielmi & Platia, 1985
[Guglielmi & Platia 1985: 210; Lodos 1998: 36; Cate 2007: 147].

Melanotus punctolineatus (Pélerin, 1829)

[Bodemeyer 1900: 116 (as *M. niger*); Guglielmi & Platia 1985: 211; Lodos 1998: 36, Lodos 1998: 36 (as *M. niger*); Cate 2007: 148].

* *Melanotus sobrinus* (Ménétriés, 1832)

[Lodos 1998: 36].

Known from Caucasus, Iran and Turkmenistan, its occurrence in Turkey is dubious.

Melanotus tenebrosus (Erichson, 1841)

[Lodos 1998: 36].

Confirmed occurrence in Turkey. Present paper → p. 33.

Melanotus turcicus Platia & Gudenzi, 2002

[Platia & Gudenzi 2002: 94; Cate 2007: 149].

* *Melanotus turkestanicus* Schwarz, 1902

[Lodos 1998: 36].

Known from Kyrgyzstan, it is improbable that the taxon occurs in Turkey.

Melanotus villosus (Fourcroy, 1785)

= *M. rufipes* (Herbst, 1784)

[Winkler 1924: 596 (as *M. rufipes*); Schenkling 1925-1927: 279, 280 (as *M. rufipes*); Horion 1953: 240; Yüksel 1970: 21, 52 (as *M. rufipes*); Guglielmi & Platia 1985: 211; Lodos 1998: 35, 36, 37, 38 (as *M. rufipes*); Lodos 1998: 36; Kesdek, Platia & Yıldırım 2006: 427; Cate 2007: 150].

Subgenus *Spheniscosomus* Schwarz, 1892

Melanotus cuneiformis (Baudi di Selve, 1871)

[Schwarz 1892: 149; Preiss & Platia 2003: 115; Cate 2007: 151].

Genus *Melanotus* species incertae sedis

* *Melanotus cranicollis* Er.

[Lodos 1998: 36].

Probably mistaken transcription of correct name.

Subfamily Negastriinae Nakane et Kishii, 1956

Genus *Negastrius* C. G.Thomson, 1859

Negastrius sabulicola (Boheman, 1851)

First record for Turkey. Present paper → p. 33.

Genus *Quasimus* Gozis, 1886

Quasimus minutissimus (Cermar, 1817)

[Bodemeyer 1900: 116 (as *Cryptohypnus*); Schenkling 1925: 216; Horion 1953: 225; Guglielmi & Platia 1985: 211; Kesdek, Platia & Yıldırım 2006: 428; Cate 2007: 191].

Genus *Zorochros* C. G.Thomson, 1859

Zorochros alysidotus (Kiesenwetter, 1858)

[Bodemeyer 1900: 116 (as *Cryptohypnus*); Guglielmi & Platia 1985: 211; Preiss & Platia 2003: 104, Kabalak & Sert 2005: 55; Cate 2007: 192].

Zorochros anatolicus Dolin & Mertlik, 2002

[Dolin & Mertlik 2002: 91; Cate 2007: 192].

Zorochros araxicola (Reitter, 1895)

[Cate 2007: 192].

Zorochros caucasicus Jagemann, 1939

[Jagemann 1939: 68; Cate 2007: 192].

27 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

Zorochros dermestoides (Herbst, 1806)

= *Z. minimus* Lacordaire, 1835

First record for Turkey. Present paper → p. 35.

Zorochros georgicus Dolin & Tschatlandze, 1980

[Cate 2007: 193].

Zorochros heyrovskyi Roubal, 1940

[Cate 2007: 193].

Zorochros kopeckyi Platia & Gudenzi, 1998

[Platia & Gudenzi 1998: 636; Cate 2007: 193].

Zorochros kurdicus Dolin & Mertlik, 2002

[Dolin & Mertlik 2002: 90; Cate 2007: 193].

Zorochros meridionalis (Laporte de Castelnau, 1840)

[Bodemeyer 1900: 116 (as *Cryptohypnus*); Horion 1953: 225; Guglielmi & Platia 1985: 211, Kabalak & Sert 2005: 55; Cate 2007: 193].

Zorochros murinus (Reitter, 1895)

[Cate 2007: 193].

Zorochros nanus (Gurjeva, 1963)

First record for Turkey. Present paper → p. 35.

Zorochros pallicrus (Desbrochers des Loges, 1875)

= *Z. delicatulus* (Schwarz, 1897)

[Cate 2007: 193].

Zorochros pilosellus (Reitter, 1895)

[Cate 2007: 194].

Zorochros quadriguttatus (Laporte de Castelnau, 1840)

First record for Turkey. Present paper → p. 35.

Zorochros quadrinaevus (Reitter, 1895)

[Cate 2007: 194].

Zorochros recellentus Dolin, 1995

[Dolin 1995: 19; Cate 2007: 194].

Zorochros stibicki (Leseigneur 1970)

[Lohse 1979: 179, Kabalak & Sert 2005: 55, Cate 2007: 194].

Subfamily Plastocerinae Crowson, 1972

= Aplastinae Stibick, 1979

Genus *Plastocerus* LeConte, 1853

Plastocerus angulosus (Germar, 1845)

[Winkler 1924: 578; Bocák 2007: 209].

Confirmed occurrence for Turkey. Present paper → p. 34.

Catalogue of the family Lissomidae Laporte, 1835

Genus *Drapetes* Dejean, 1821

Drapetes flavipes subparalellus Pic, 1908

[Pic 1908: ?; Smetana 2007: 207].

Drapetes mordelloides (Host, 1789)

= *D. biguttatus* (Piller & Mitterpacher, 1783)
[Reitter 1921: 74; Smetana 2007: 207].

Catalogue of the family Melasidae Leach, 1817

= Melasidae Fleming, 1821
= Eucnemidae Eschscholz, 1829
= Phylloceridae Reitter, 1905

The classification follows Muona (2007)

Subfamily Macraulacinae Fleutiaux, 1922

Tribe **Euryptychini** Mamaev, 1976

Genus *Dromaeolus* Kiesenwetter, 1858

Dromaeolus barnabita (A. Villa & J. B. Villa, 1838)

[Mertlik, Jeniš & Zbuzek 2007: 92].

Dromaeolus maronita Bonvouloir, 1871

[Reitter 1921: 78 (as *Megathambus*); Winkler 1924: 618 (as *Megathambus*)].

Subfamily Melasinae Leach, 1817

Tribe **Dirhagini** Reitter, 1911

Genus *Farsus* Jacquelin du Val, 1860

Farsus dubius (Piller & Mitterpacher, 1783)

First records for Turkey. Present paper → p. 35.

Genus *Microrhagus* Dejean, 1833

Microrhagus emyi (Rouget, 1856)

First records for Turkey. Present paper → p. 35.

Genus *Rhacopus* Hampe, 1855

Rhacopus sahlbergi (Mannerheim, 1823)

29 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

[Mertlik, Jeniš & Zbuzek 2007: 93].

Genus *Epiphanis* Eschscholtz, 1829

Epiphanis cornutus Eschscholtz, 1829

[Mertlik, Jeniš & Zbuzek 2007: 93].

Tribe **Melasini** Fleming, 1821

Genus *Isorhipis* Lacordaire, 1835

Isorhipis melasoides (Laporte, 1835)

[Muona 2007: 84].

Isorhipis nigriceps (Mannerheim, 1823)

[Reitter 1921: 77; Winkler 1924: 617; Muona 2007: 84].

Genus *Melasis* A. G. Olivier, 1790

Melasis buprestoides (Linnaeus, 1761)

[Muona 2007: 84].

Subfamily Phyllocerinae Reitter, 1905

Tribe **Anelastini** Reitter, 1921

Genus *Anelastes* Kirby, 1819

Anelastes barbarus P. H. Lucas, 1846

[Muona 2007: 81].

Tribe **Phyllocerini** Reitter, 1905

Genus *Phyllocerus* Le Peletier de Saint-Fargeau & Audinet-Serville, 1825

Phyllocerus diversicornis Pic, 1905

[Reitter 1921: 89; Winkler 1924: 578; Wurst 1996: 55; Muona 2007: 81].

Phyllocerus oreadis Wurst, 1996

[Wurst 1996: 52; Muona 2007: 81].

Catalogue of the family Throscidae Laporte, 1840

The classification follows Leseigneur (2007)

Subfamily Throscinae Laporte, 1840

Tribe Throscini Laporte, 1840

Genus *Aulonothroscus* Horn, 1890

Aulonothroscus brevicollis (Bonvouloir, 1859)

[Leseigneur 2007: 87].

Aulonothroscus maximus (Pic, 1938)

[Leseigneur 2007: 88].

Genus *Trixagus* Kugelann, 1794

Trixagus atticus Reitter, 1921

First record for Turkey. Present paper → p. 36.

Trixagus carinifrons (Bonvouloir, 1859)

[Leseigneur 2007: 88].

Trixagus dermestoides (Linnaeus, 1767)

[Leseigneur 2007: 88].

Trixagus duvalii (Bonvouloir, 1859)

First record for Turkey. Present paper → p. 36.

Trixagus elateroides (Heer, 1841)

[Leseigneur 2007: 88].

Trixagus exul (Bonvouloir, 1859)

[Leseigneur 2007: 88].

Trixagus gracilis Wollaston, 1854

[Leseigneur 2007: 88].

Trixagus obtusus (Curtis, 1827)

[Reitter 1921: 72; Leseigneur 2007: 89].

Trixagus orientalis (Bonvouloir, 1859)

[Leseigneur 2007: 89].

New distributional data of the family Elateridae, Melasidae and Throscidae

Family Elateridae Leach, 1815

***Adrastus samedovi* Dolin & Agajev, 1974**

Specimens studied. Turkey, prov. Kars, 3 km W of Kağızman, Aras river, 29.vi.2006, 1 ♂, J. Mertlik leg. et coll., G. Platia det. 2007; Turkey, prov. Erzurum, 10 km NW of İspir, 24.vi.2006, 1 ♂, J. Mertlik leg. et coll., G. Platia det. 2007.

Distribution. Azerbaijan, Iran (Cate 2007).

First records for Turkey.

***Agriotes lapicida* (Falderman, 1835)**

Specimen studied. Turkey, prov. Malatya, 10 km SE of Yukan Ulupinar, 30 km SE Derende, 26.- 27.v.2001, 1 ♀, P. Kabátek leg., J. Mertlik coll., G. Platia det. 2007.

Distribution. Azerbaijan, Armenia, Georgia, Russia: South European Territory, Iran (Cate 2007); Turkey (Lodos 1998),

Confirmed occurrence in Turkey.

***Agriotes meticulosus* Candèze, 1863**

Specimen studied. Turkey, prov. Erzurum, Çobandede (Köprüköy), Aras Nehri, 13.- 15.vii.1997, 1 ♀, J. Mertlik leg. et coll.

Distribution. Azerbaijan, Armenia, Georgia, Russia: South European Territory, Afghanistan, Iran, Iraq, Kazakhstan, Mongolia, Pakistan, Saudi Arabia, Syria, Tajikistan, Turkmenistan, Uzbekistan (Cate 2007).

First record for Turkey.

***Ampedus rufipennis* (Stephens 1830)**

Specimens studied. Turkey, prov. Bursa, Uludag 1150m, 5.vi.2000, 1 ♂, V. Skoupý leg., J. Mertlik coll.; Turkey, prov. Antalya, Termessos, 3.v.1992, H. Schmid leg., G. Platia et P. Cate coll. (Wien, Austria).

Distribution. Azerbaijan, Armenia, Austria, Bosnia Herzegovina, Bulgaria, Croatia, Czech Republic, Denmark, France, Great Britain, Germany, Georgia, Greece, Hungary, Italy, Moldavia, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Russia: South European Territory, Sweden, Ukraine, Serbia and Montenegro, Algeria, Morocco, Iran (Cate 2007).

First records for Turkey.

***Apotarsus angustulus* (Kiesenwetter, 1858)**

Specimen studied. Turkey, prov. Kars, Susuz env., 6.vi.1994, 1 ♂, V. Skoupý leg., J. Mertlik coll.

Distribution. Azerbaijan, Austria, Belgium, Bulgaria, Czech Republic, France, Great Britain, Germany, Poland, Romania, Slovakia, Russia: South European Territory, Switzerland, Ukraine (Cate 2007).

First record for Turkey.

***Denticollis rubens* Piller et Mitterpacher, 1783**

Specimen studied. Turkey, prov. Artvin, Cankurtaran Geçidi, 20 km E Hopa, 700 m, 4123N 4133E, 26.-27.vi.1999, 1 ♂, M. Rejzek leg. et coll. (Norwich, UK), J. Mertlik det.

Distribution. Austria, Belgium, Bosnia Herzegovina, Bulgaria, Belarus, Croatia, Russia: Central European Territory, Czech Republic, Denmark, Estonia, France, Germany, Georgia, Hungary, Italy, Latvia, Liechtenstein, Luxembourg, Moldavia, Norway, Poland, Romania, Serbia and Montenegro, Slovakia, Slovenia, Russia: South European Territory, Sweden, Switzerland, Ukraine (Cate 2007), Spain (Molino-Olmedo 2005).

First record for Turkey.

***Elathous buyssoni* Reitter, 1890**

Specimen studied. Turkey, prov. Artvin, Barhal-Yayla, SW Sarigol, 1900-2500 m, 7.viii.1978, 1 ♀, M. Zeising leg., G. Platia det. et coll.; Turkey, prov. Amasya, 22.viii.1990, 1 ♀, S. Dacatra leg., G. Platia det. et coll.

Distribution. Azerbaijan, Armenia (Cate 2007).

First record for Turkey.

***Elater ferrugineus ferrugineus* Linnaeus, 1758**

Specimen studied. Turkey, prov. İçel (Mersin), Gülnar, 50 km N Kuseçobali, Tasdustu 16 W, 1480m, 26.vii.2006, 1 ♂ (f. *morio* Schilsky, 1888), on *Quercus* sp.; M. Coskun & N. Jansson leg., N. Jansson coll. (Sweden), G. Platia det.

Distribution. Albania, Armenia, Austria, Belgium, Bosnia Herzegovina, Bulgaria, Russia (Central European Territory), Czech Republic, Denmark, France, Great Britain, Germany, Georgia, Greece, Hungary, Italy, Moldavia, The Netherlands, Romania, Slovakia, Spain, Russia (South European Territory), Sweden, Switzerland, Ukraine (Cate 2007).

Confirmed occurrence in Turkey.

***Hemicrepidius niger* (Linnaeus, 1758)**

Specimen studied. Turkey, prov. Tekirdağ, Çorlu, 31.v.1986, 1 ♂, Kadlec & Voříšek leg., J. Mertlik coll.

Distribution. Azerbaijan, Albania, Austria, Belgium, Bulgaria, Belarus, Croatia, Russia (Central European Territory), Czech Republic, Denmark, Estonia, Finland, France, Germany, Georgia, Hungary, Italy, Latvia, Liechtenstein, Lithuania, Luxembourg, The Netherlands, Norway, Russia (North European Territory), Poland, Romania, Slovenia (Cate 2007), Slovakia (Laibner 2000); Turkey (Kabalak & Sert 2005).

Confirmed occurrence in Turkey.

***Ischnodes sanguinicollis* (Panzer, 1793)**

Specimens studied. Turkey, prov. İçel, Erdemli env., 8 km NW of Arslanlı, 24.-26.v.1995, 1 ♂, 1 ♀, J. Mertlik leg. et coll.; Turkey, prov. Adana/Gaziantep, Nurdağı Geçidi 1200m, 1.- 4.vi.1992, 1 ♂, M. Bednářík leg., J. Mertlik coll.; Turkey, prov. Muş, Buğlan Geçidi 1600m, 14.- 15.vi.1992, 1 ♂, M. Bednářík leg., J. Mertlik coll.; Turkey, prov. Manisa, 40 km S of Balikesir, former oaken forest-clear felling, 21.vi.2002, more larvae, in cavity of stump of oak, J. Mertlik leg. et coll.; Turkey, prov. Isparta, Davraz Dağı, Yukarıgökdere env., pasture woodland, 23.v.2001, more larvae, in cavity of stump of oak, J. Mertlik leg. et coll.

Distribution. Azerbaijan, Austria, Belgium, Bosnia Herzegovina, Croatia, Czech Republic, Denmark, France, Great Britain, Germany, Greece, Hungary, Italy, Moldavia, Poland, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Ukraine, Iran, Japan, Syria (Cate 2007); Turkey (Platia 1994).

Confirmed occurrence in Turkey.

33 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

***Isidus letourneuxi* Pic, 1902**

Specimen studied. Turkey, prov. Antalya, Alanya, 16.vi.2006, 1 ♂, L. Ernest leg., J. Mertlik coll.

Distribution. Egypt (Cate 2007).

First record for Turkey.

***Lacon unicolor* (Candèze, 1874)**

Specimen studied. Turkey, prov. Mardin, Haberli, 33 km SE Midyat (3717N 4135E), 17.v.2001, 1 ♀, S. Kadlec. leg. et coll. (Litvínov, Czech Republic), J. Mertlik det. 2007.

Distribution. Iran (Cate 2007).

First record for Turkey.

***Melanotus brignolii* Guglielmi & Platia, 1985**

Specimens studied. Turkey, prov. Antalya, S Taurus, 30 km NW Alanya, 1200m, 14.- 28.vii.2001, 2 ♂♂, M. Hiermeier leg., J. Mertlik coll., G. Platia det. 2004; Turkey, prov. Antalya, near İncekum, vii.2000, 1 ♂, Lizler et Werner leg., G. Platia coll.

Distribution. Greece (Cate 2007).

First records for Turkey.

***Melanotus fulvus* Reitter, 1891**

Specimen studied. vil. Kars, 3 km W of Kağızman, Aras river, 29. vi. 2006, 1 ♂, Bořivoj Zbuzek leg. et coll., J. Mertlik det. 2007.

Distribution. Iran, Iraq, Turkmenistan (Cate 2007).

First record for Turkey.

***Melanotus tenebrosus* (Erichson, 1841)**

Specimen studied. Turkey, prov. Artvin, 35 km W Sarigol, 5 km NW Yaylalar, 1.vii.2003, 1 ♂, E. et P. Hajdaj leg., J. Mertlik coll.

Distribution. Armenia, Austria, Bosnia Herzegovina, Croatia, Czech Republic, France, Georgia, Greece, Hungary, Italy, Portugal, Romania, Slovakia, Slovenia, Spain, Russia: South European Territory, Switzerland (Cate 2007).

Confirmed occurrence for Turkey.

***Negastrius sabulicola* (Bohemian, 1851)**

Specimen studied. Turkey, prov. Bursa, Ulubat Gölü, 27.v.1996, 1 ♂, Z. Andrš leg., J. Mertlik coll.

Distribution. Austria, Belgium, ?Bosnia Herzegovina, Bulgaria, Belarus, Russia: Central European Territory, Czech Republic, Denmark, France, Great Britain, Germany, Greece, Hungary, Italy, Latvia, Lithuania, The Netherlands, Russia: North European Territory, Poland, Romania, Slovakia, Sweden, Ukraine (Cate 2007).

First record for Turkey.

***Paracardiophorus musculus* (Erichson, 1840)**

Specimens studied. Turkey, prov. Sivas, 10 km SE of Koyulhisar, 11.vi.2000, 1 ♂, J. Mertlik leg. et coll.; Turkey, prov. Artvin, 5 km W of Şavşat, Okçular river, 28.vi.2006, 2 ♂♂, J. Mertlik leg. et coll.; SW Turkey, Taurus,

[Liciae], ex Winkler coll., 1 ♂, G. Platia coll.

Distribution. Austria, Bulgaria, Russia: Central European Territory, Czech Republic, France, Germany, Georgia, Greece, Hungary, Italy, Lithuania, Poland, Romania, Slovakia, Spain, Russia: South European Territory, Switzerland, Ukraine, Algeria, Tunisia, Russia: East Siberia, Russia: west Siberia (Cate 2007).

First records for Turkey.

Plastocerus angulosus (Germar, 1844)

Specimens studied. Turkey, prov. Çanakkale, Bahramkale (Assos), 19.- 24.vii.1991, 2 ♂♂, Vít Kubáň leg., J. Mertlik coll.; Turkey, prov. İçel (Mersin), 15 km W of Mersin, 1.- 5.vii.1992, 3 ♂♂, D. Hauck leg., J. Mertlik coll.; Turkey, prov. Adana, Pozanti env., Ak Dağı, 10.- 12.vii.1997, 8 ♂♂, J. Mertlik leg.; Turkey, prov. Isparta, Davraz Dağı, Yukarıgökdere env., 12.- 13.vii.2000, 1 ♂, R. Lízler et K. Werner leg., J. Mertlik coll.; Turkey, prov. Isparta, Isparta, Davraz Dağ, 25.v.2001, 1 ♂, M. Snížek leg., G. Platia coll.; Turkey, prov. Eskişehir, Eskisehir env., Porsuk barajı lake, 7.vii.1993, 1 ♂, A. Šlachta leg., J. Mertlik coll.; Turkey, prov. Antalya, Beldibi p. Kemeri, Bey Dağları, 1.- 15.vii.1998, 4 ♂♂, P. Bulirsch leg., J. Mertlik coll.; Turkey, prov. Adana, Cinarlı, 30.vi.1997, 1 ♂, J. Hájek leg., J. Mertlik coll.; Turkey, prov. Eskişehir, Alpu, 5.vii.1996, 1 ♂, J. Hájek leg., J. Mertlik coll.; Turkey, prov. Bursa, Uludağ, 20 km S of Bursa, Çaybaşı env., 4.vii.2006, 2 ♂♂, J. Mertlik leg. et coll.; Turkey, prov. Adana, Osmaniye, Karatepe, 3.- 7.vii.1994, 3 ♂♂, 1 ♀, P. Kabátek leg., J. Mertlik coll.; Turkey, prov. Izmir, Zeytindağ, 19.vi.2002, 1 ♀ (f. *atricolor* Reitter, 1896), M. Samek leg. et coll. (Skalice, Czech Republic); Turkey, prov. Zonguldak, Karabük, 20 km E, 22.- 23.vi.1996, 2 ♂, P. Zahradník leg. et coll., J. Vávra det. 2006, J. Mertlik rev. 2007; Turkey, prov. İçel (Mersin), Namrun, 26.vii.1985, 1 ♂, Niehuis leg., G. Platia coll.

Distribution. Turkey (Bocák 2007).

Confirmed occurrence for Turkey.

Procræterus tibialis (Lacordaire, 1835)

Specimen studied. Turkey, prov. Edirne, Edirne, 16.v.1993, 1 ♂, M. Šárovec leg., J. Mertlik coll.

Distribution. Austria, Belgium, Bosnia Herzegovina, Bulgaria, Croatia, Russia: Central European Territory, Czech Republic, Denmark, Finland, France, Great Britain, Germany, Greece, Hungary, Italy, Lithuania, ?Luxembourg, Moldavia, Norway, Poland, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Ukraine, Serbia and Montenegro (Cate 2007).

First record for Turkey.

Selatosomus ampliformis Reitter, 1910

Specimen studied. Turkey, prov. Kars, Digor env., pass 2130m, 30 km SE of Kars, 12.- 13.vi.1999, 2 ♀♀, T. Lackner leg., V. Dušánek (Zábřeh, Czech Republic) et J. Mertlik coll.

Distribution. Azerbaijan, Armenia (Cate 2007).

First record for Turkey.

Stenagostus rosti (Schwarz, 1897)

Specimen studied. Turkey, prov. Artvin, 15 km W of Borçka, Cankurtaran Geçidi, 28.vi.2003, 1 ♀, E. et P. Hajdaj leg., J. Mertlik coll.

Distribution. Azerbaijan, Georgia, Russia: South European Territory (Cate 2007); Turkey (Orlov 1989, Lodos 1998).

Confirmed occurrence in Turkey.

***Zorochros dermestoides* (Herbst, 1806)**

Specimens studied. Turkey, prov. Bolu, Abant gölü lake, 1200 m, 4.vi.1996, 1 ex. J. Vávra leg. et coll. (Ostrava, Czech Republic), J. Mertlik det.; 23.-24.vi.1996, 4 ex., J. Vávra leg. et coll., J. Mertlik det.; Turkey, prov. Kastamonu/Çankırı, Ilgaz Dağ, 3.vi.1991, 1 ♀, V. Skoupý leg., J. Mertlik coll.; Turkey, prov. Erzurum, 12 km E of Tortum, 30.vi.2006, 1 ♀, J. Mertlik leg. et coll.

Distribution. Andorra, Armenia, Austria, Belgium, Bosnia Herzegovina, Bulgaria, Belarus, Croatia, Russia: Central European Territory, Czech Republic, Denmark, Estonia, Finland, France, Germany, Great Britain, Georgia, Greece, Ireland, Italy, Lithuania, Macedonia, Moldavia, The Netherlands, Norway, Russia: North European Territory, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Russia: South European Territory, Sweden, Switzerland, Ukraine, Serbia and Montenegro, Morocco (Cate 2007); Hungary (Merkl & Mertlik 2005).

First records for Turkey.

***Zorochros nanus* (Gurjeva, 1963)**

Specimens studied. Turkey, prov. Iskenderun, Tekebasi p. Samandağı, Asi Nehri, 29.vi.1992, 1 ex., P. Bulirsch leg., J. Mertlik coll., W. Dolin det. 1996; Turkey, prov. Iskenderun, Ulucinar env., 11.- 12.iv.1992, 2 ex., O. Hovorka leg., J. Mertlik coll.; Turkey, prov. Iskenderun, Hatay, Nur Dağları Mts, Dorytoi, 3.v.2005, more ex., Z. Malinka leg., G. Platia & Z. Malinka coll. (Opava, Czech Republic).

Distribution. Kyrgyzstan (Cate 2007).

First records for Turkey.

***Zorochros quadriguttatus* (Laporte de Castelnau, 1840)**

Specimens studied. Turkey, prov. Bursa, Iznik env., 22.vi.1992, 2 ex., P. Bulirsch leg., J. Mertlik coll.; Turkey, prov. Trabzon, 10 km S of Araklı, Küçükdere env., 23.vi.2006, 3 ex., J. Mertlik leg. et coll.

Distribution. Armenia, Austria, Bulgaria, Czech Republic, Greece, Hungary, Italy, Macedonia, Poland, Portugal, Romania, Slovakia, Spain, Russia: South European Territory (Cate 2007).

First records for Turkey.

Family Melasidae Leach, 1817

***Farsus dubius* (Piller & Mitterpacher, 1783)**

Specimens studied. Turkey, prov. Isparta, Yukarıgökdere env., Davraz Dağı, 18.- 23.vii.1998, 1 ex., R. Lízler & K. Werner leg., J. Mertlik coll.; Turkey, prov. Isparta, Yukarıgökdere env., Davraz Dağı, vii.1999, 8 ex., K. Werner leg., J. Mertlik coll.; Turkey, prov. Adana, Kozan env., vii.1998, 1 ex., K. Werner leg., J. Mertlik coll.; Turkey, prov. İçel, Erdemli env., 8 km NW of Arslanlı, 24.- 26.v.1995, 1 ex., J. Mertlik leg. et coll.; Turkey, prov. İçel, Erdemli env., 8 km NW of Arslanlı, 24.v.1998, 1 ex., J. Mertlik leg. et coll.

Distribution. Austria, Czech Republic, France, Hungary, Italy, Romania, Spain, Russia (South European Territory), “Yugoslavia” (Muona 2007).

First records for Turkey.

***Microrhagus emyi* (Rouget, 1856)**

Specimen studied. Turkey, prov. Zonguldak, Karabük, 20 km E, 22.- 23.vi.1996, 2 ♂, P. Zahradník leg. et coll. (Mníšek pod Brdy, Czech Republic), J. Vávra det. 2006, J. Mertlik rev. 2007.

Distribution. Austria, Croatia, Czech Republic, France, Italy, Germany, Hungary, Poland, Romania, Russia: Central European Territory, Slovakia, Spain, Sweden, Switzerland, Ukraine, „Yugoslavia“ (Muona 2007); Slovenia,

Bulgaria, Greece and Russia: South European Territory (Mertlik, Jeniš & Zbuzek 2007).

First record for Turkey.

Family Throscidae Laporte, 1840

Trixagus atticus Reitter, 1921

Specimens studied. Turkey, prov. Zonguldak, Karabük, 20 km E, 22.- 23.vi.1996, 1 ♂, P. Zahradník leg. et coll., J. Mertlik det. 2007; Turkey, prov. Zonguldak, Karabük, 20 km E, 27.vi.2002, 1 ♀, J. Mertlik leg. et coll., L. Leseigneur det. 2004, J. Mertlik rev. 2007.

Distribution. Denmark, Finland, France, Germany, Greece, Italy, Romania, Russia (South European Territory), Sweden, Switzerland, Afghanistan, Cyprus, Turkmenistan (Leseigneur 2007).

First records for Turkey.

Trixagus duvalii (Bonvouloir, 1859)

Specimens studied. Turkey, prov. Konya, Beyşehir env., Yeşildağ, 29.v.1995, 1 ♀, T. Kopecký leg., L. Leseigneur det. 2000, J. Mertlik coll.; Turkey, prov. Adiyaman, Azaplı Gölü lake, Gölbaşı env., 17.- 18.v.2005, 6 ♂♂, 3 ♀♀, Z. Malinka leg. et coll. (Opava, Czech Republic) (1 ♂, 1 ♀, J. Mertlik coll.).

Distribution. Austria, Czech Republic, Denmark, Finland, France, Great Britain, Germany, Greece, Hungary, Italy, Poland, Romania, Slovakia, Slovenia, Spain, Russia (South European Territory), Sweden, Cyprus (Leseigneur 2007).

First records for Turkey.

Acknowledgements

The authors thank Lucien Leseigneur (Meylan, France) for invaluable advice and reviewing the manuscript. It is a great pleasure to express our sincerest gratitude to Nilay Gülpereçin and (Natural History Application and Research Center, Natural History Museum, University of Ege, Bornova, Izmir, Turkey), Serdar Tezcan (Department of Plant Protection, Faculty of Agriculture, University of Ege, Bornova, Izmir, Turkey) and Mahmut Kabalak (Hacettepe University Faculty of Science, Department of Biology, Beytepe, Ankara, Turkey) for their generously allowing us to study the Turkish literature. We also thank Jiří Hájek (National Museum, Department of Entomology, Prague, Czech Republic) for being allowed to examine type material in its care.

Our gratitude is also due Zdeněk Andrš (Černošín, Czech Republic), Michal Bednářík (Olomouc, Czech Republic), Stanislav Benedikt (Plzeň, Czech Republic), Petr Bulirsch (Lovosice, Czech Republic), Václav Dušánek (Zábřeh, Czech Republic), Ladislav Ernest (Nymburk, Czech Republic), Evžen Hajdaj & Patrik Hajdaj (Ježov, Czech Republic); Oldřich Hovorka (Dobříš, Czech Republic), Nicklas Jansson (Linköping, Sweden), Petr Kabátek (Praha, Czech Republic), Vít Kubáň (Brno, Czech Republic), Stanislav Kadlec (Litvínov, Czech Republic), Tomáš Kopecký (Hradec Králové, Czech Republic), Tomáš Lackner (Amsterdam, The Netherlands), Robert Lízler (Hradec Králové, Czech Republic), Roman Lohaj (Košice, Slovakia), Zdeněk Malinka (Opava, Czech Republic), Martin Samek (Skalice, Czech Republic), Vladimír Skoupý (Žilina, Czech Republic), Milan Šárovec (Pardubice, Czech Republic), Jiří Vávra (Ostrava, Czech Republic), †Karl Werner (Peiting, Germany) Petr Zahradník (Mníšek pod Brdy, Czech Republic), Bořivoj Zbuzek (Praha, Czech Republic) for lending us material for this study.

References

- BOCÁK L. 2007: Family Plastoceridae, 209. In LÖBL I. & SMETANA A. (eds.): *Catalogue of Palaearctic Coleoptera, Vol. 4. Elateroidea - Derodontoidea - Bostrichoidea - Lymexyloidea - Cleroidea - Cucujoidea*. Apollo Books, Stenstrup, 935 pp.
- BODEMEYER E. V. 1900: Quer durch Kleinasien in den Bulghan Dagh. - Die Druck und Verlags-Aktiengesellschaft. *Vormals Dölter*, Emmedingen, 169 pp.
- BOOTH R.G., COX M.L. & MADGE R.B. 1990: In: Guides to Insects of Importance to Man, 3. Coleoptera. *International Institute of Entomology*, 384 pp.
- BUYSSON H. DU 1891: Espèces nouvelles d'Elaterides. *L'Echange Revue Linnaéenne*, 7, 133-136.
- CATE P. & PLATIA G. 1989: A new species of the genus *Cidnopus* Thomson from Greece: *Cidnopus macedonicus* n. sp. (Elateridae, Coleoptera). *Koleopt. Rdsch.*, 59: 185-188.
- CATE P. & PLATIA G. 1997: A new species of *Agriotes* Eschscholtz (Coleoptera, Elateridae) from Greece, Turkey and Syria. *Z. Arb. Gem. Öst. Ent.*, 49: 109-113.
- CATE P. C., PLATIA G. & SCHIMMEL R. 2002: New species and records of click beetles (Coleoptera: Elateridae) from Iran, with a checklist of known species. *Folia Heyrovskiana*, 10 (1): 25-68.
- CATE P. G. 2007: Family Elateridae, 89-209. In LÖBL I. & SMETANA A. (eds.): *Catalogue of Palaearctic Coleoptera, Vol. 4. Elateroidea - Derodontoidea - Bostrichoidea - Lymexyloidea - Cleroidea - Cucujoidea*. Apollo Books, Stenstrup, 935 pp.
- DOLIN V. G. 1995: Neue *Zorochrus*-Arten (Coleoptera, Elateridae) aus dem Transkaukasus und Zentral-Asien. *Zeitschrift der Arbeitsgemeinschaft Österreichischer Entomologen*, 47 (1/2): 19-24.
- DOLIN V. G. & MERTLIK J. 2002: Drei neue Klein-Asiatischen und Iranischen *Zorochrus*-Arten (Coleoptera, Elateridae, Negastriinae). *Vestnik zoologii*, 36 (6): 89-93.
- DUŠÁNEK V. & MERTLIK J. 2007: Kraj Athousů - aneb běžný den entomologů v Turecku (Coleoptera, Elateridae) [Land of Athouses - the usual day of entomologists in Turkey (Coleoptera, Elateridae)]. *Elateridarium*, 1: 68-72. (in Czech).
- FAIRMAIRE M.L. 1866: Notice sur les Coleopteres recoltes par M.J. LEDERER (sur le Bosz-Dagh, Asie Mineure). *Annales de la Societe Entomologique de France*, 35: 249-280.
- GÜL-ZÜMREOGLU S. 1972: *Izmir Bölge Zirai Mücadele Araştırmaları Böcek ve Genel Zararlilar Katalogu*, 1928-1969, 119 pp.
- GUGLIELMI A. & PLATIA G. 1985: Contributo alla conoscenza degli Elateridi di Grecia e Turchia (Coleoptera). *Fragm. Entomol.*, 18 (1): 169-224.
- GURJEVA E. L. 1972: A review of Palearctic species of the genus *Agriotes* Esch. (Col. Elateridae). *Ent. Obzr.*, 51 (4): 859-877.
- GURJEVA E. L. 1989: Fauna SSSR. Elateridae subf. Athoinae (Tribe Ctenicerini). *Nauka*, Leningrad, 12 (3), 293 pp.
- HEYDEN L. V. O., REITTER E. & WEISE J. 1906: *Catalogus Coleopterorum Europae, Caucasi et Armeniae Rossiae*. Berlin, 774 pp.
- HORION A. 1953: Faunistik der mitteleuropäischen Käfer. *Entom. Arb. Mus. Frey; Sonderband*. München.- 340 pp.
- CHASSAIN J. 1998: Note sur le Genre *Limoniscus* avec désignation d'un Lectotype et description d'une espèce nouvelle (Coleoptera, Elateridae). *Le Coleoptériste*, 33: 119-122.
- JAGEMANN E. 1939: Einige neue Elateriden-Formen aus dem paläarkt. Faunengebiet. *Entomologisches Nachrichtenblatt*, 13 (1/2): 68-70.
- KABALAK M. & SERT O. 2005: Ankara İli Elateridae (Coleoptera) familyası türleri üzerinde faunistik çalışmalar (Faunistical studies on species of the family Elateridae (Coleoptera) in Ankara). *Türkiye entomoloji dergisi*, 29 (1): 49-60. (In Turkish)
- KABALAK M. & SERT O. 2006: SCIENTIFIC NOTE. New record of *Melanotus bajulus* (Erichson, 1841) (Coleoptera: Elateridae) for the Turkish fauna. *The Coleopterists Bulletin*, 60 (3): 216.
- KESDEK M., PLATIA G. & YILDIRIM E. 2006: Contribution to the Knowledge of Click-Beetles (Coleoptera: Elateridae) Fauna of Turkey. *Entomofauna*, 27 (35): 417-432.
- KISHII T. 1999: A check-list of the family Elateridae from Japan (Coleoptera). *Bulletin of the Heian High School*, Kyoto, 42: 1-144.
- KUBÁŇ V. 1995: Towards the occurrence of click-beetles *Ampedus (Brachygonus) megerlei*, *Ampedus (Brachygonus) dubius* and *Ectamenogonus montandoni* (Coleoptera: Elateridae) on the territory of the Czech Republic and Slovak Republic. *Klapalekiana*, 31 (1-2): 23-30.

- LAIBNER S. 2000: Elateridae of the Czech and Slovak Republics České a Slovenské republiky. *Kabourek*, Zlín: 292 p. (In Engl. & Czech).
- LAWRENCE J.F., HASTINGS A.M., DALLWITZ M.J., PAINE T.A. & ZURCHER E.J. 2000 onwards. Elateriformia (Coleoptera): descriptions, illustrations, identification, and information retrieval for families and subfamilies. Version: 9th October 2005. <http://delta-intkey.com>
- LESEIGNEUR L. 2007: Family Throscidae, pp. 87-89. In LÖBL I. & SMETANA A. (eds.): *Catalogue of Palaearctic Coleoptera, Vol. 4. Elateroidea - Derodontoidea - Bostrichoidea - Lymexyloidea - Cleroidea - Cucujoidea*. Apollo Books, Stenstrup, 935 pp.
- LODOS N. 1998: Türkiye Entomolojisi VI (Genel, Uygulamalı ve Faunistik). *Ege Üniversitesi Zirarat Fakültesi Yayınları* No: 529, 300 pp.
- LOHSE G. A. 1979: Elateridae. In Freude H., Harde K. W., Lohse G. A.: Die Käfer Mitteleuropas. 6. *Goecke & Evers*, Krefeld, 103-186.
- MERTLIK J. 2000: Contribution to the knowledge of Turkish species of the genus *Selatosomus* (Coleoptera: Elateridae). *Folia Heyrovskiana*, 8 (1): 1-15.
- MERTLIK J. 2005: Description of three new species of family Elateridae (Coleoptera from Syria and Turkey. *Folia Heyrovskiana*, 12 (4): 167-173.
- MERTLIK J. & DUŠÁNEK V. 2006: Description of five new species of click-beetles (Coleoptera, Elateridae) from the Palaearctic region with remarks about the distributions 22 additional species. *Folia Heyrovskiana*, 13 (4): 145-162.
- MERTLIK J., JENIŠ I. & ZBUZEK B. 2007: New records on the distribution of some species of the family Eucnemidae (Coleoptera). *Elateridarium*, 1: 92-96.
- MUONA J. 2007: Family Eucnemidae, pp. 81-86. In LÖBL I. & SMETANA A. (eds.): *Catalogue of Palaearctic Coleoptera, Vol. 4. Elateroidea - Derodontoidea - Bostrichoidea - Lymexyloidea - Cleroidea - Cucujoidea*. Apollo Books, Stenstrup, 935 pp.
- ORLOV V. N. 1989: Larvae of two species of Elateridae beetles (Coleoptera, Elateridae) from the Caucasus. *Rev. D'Ent. URSS*, 67 (2): 329-332.
- PENTHER A. & ZEDERBAUER E. 1905: Ergebisse einer naturwissenschaftlichen Reise zum Erdschias-Dagh (Kleinasien). *Annal. Naturh. Mus. Wien*, 20 (3): 99-290.
- PIC M. 1903: Captures notes diverses et diagnoses. *Echange*, 19: 129..
- PLATIA G. 1989: Descrizione di nuove specie di Elateridi della Grecia e Turchia (Coleoptera, Elateridae). *Lavori Soc. Ven. Sc. Nat.*, 14 (1): 79-86.
- PLATIA G. 1994: Fauna d'Italia, Coleoptera Elateridae. *Edizioni Calderini*, Bologna, 429 pp.
- PLATIA G. 2001: Description de trois nouvelles espèces du genre *Ampedus* Dejean de la région palearctique, avec une synonymie nouvelle (Coleoptera: Elateridae). *Biocosme Mésogéen*, Nice, 18 (2): 65-73.
- PLATIA G. 2003: Descrizione di otto nuove di *Agriotes* del gruppo nuceus Fairmaire della Turchia con chiave di determinazione all' specie note (Insecta, Coleoptera, Elateridae). *Il Naturalista Valtellinese - Atti Mus. Civ. Stor. Nat. Morbegno*, 14: 13-28.
- PLATIA G. 2003: Descrizione di nuove specie di *Athous* Eschscholtz del sottogenere *Orthathous* Reitter della Turchia (Coleoptera, Elateridae). *Lambillionea*, CIII, 2: 263-268.
- PLATIA G. 2004: Nuovi Elateridi d'Europa, Turchia e Quatar con note tassonomiche e genemiche su alcune specie (Coleoptera: Elateridae). *Bol. S.E.A.*, 35: 39-45.
- PLATIA G. 2004: Descrizione di nuovi Elateridi della regione Palearctica con note geonomiche e sinonimiche. (Coleoptera, Elateridae). *Lambillionea*, 104 (1): 98-108.
- PLATIA G. & CATE P. 1990: Note sistematiche e sinonimiche su Elateridi Paleartici (Coleoptera Elateridae). *Boll. Soc. ent. ital.*, Genova, 122 (2): 111-114.
- PLATIA G. & GUDENZI I. 1996: Descrizione di tre nuove specie di Elateridi di Grecia, Turchia ed Isola di Cipro con note geonomiche e sinonimiche (Coleoptera, Elateridae). *Boll. Mus. reg. Sci. nat.*, Torino, 14 (1): 139-150.
- PLATIA G. & GUDENZI I. 1996: Descrizione di tre nuove specie di *Athous* Eschscholtz del sottogenere *Ortathous* Reitter della Turchia con chiave di determinazione dei maschi delle specie note (Coleoptera, Elateridae). *Lambillionea*, 96 (1): 235-241.
- PLATIA G. & GUDENZI I. 1998: Note tassonomiche e faunistiche su Elateridi del Vicino Oriente (Coleoptera, Elateridae). *Bollettino dell'Associazione Romana di Entomologia*, 53 (1-4): 49-62.
- PLATIA G. & GUDENZI I. 1998: Descrizione di nuovi Elateridi di Grecia e Turchia con note e chiave di determinazione per gli *Agriotes* del gruppo *gurgistanus* Faldermann. (Coleoptera, Elateridae). *Lambillionea*, 98 (4): 629-640.

39 Catalogue of the family Cebrionidae, Elateridae, Lissomidae, Melasidae and Throscidae from Turkey

- PLATIA G. & GUDENZI I. 1999: Descrizione di nuove specie di Elateridi della regione Paleartica con note geonomiche e sinonimiche (Insecta Coleoptera Elateridae). *Quaderno di Studi e Notizie di Storia Naturale della Romagna*, 11: 17-31.
- PLATIA G. & GUDENZI I. 2000: Descrizione di un nuovo genere e tredici nuove specie di Elateridi della regione paleartica con note geonomiche (Insecta Coleoptera Elateridae). *Quad. Studi Nat. Romagna*, 13, suppl.: 65-83.
- PLATIA G. & GUDENZI I. 2000: Revisione dei *Cardiophorus* Eschscholtz della Turchia. I: Le specie bicolori. (Coleoptera, Elateridae, Cardiophorinae). *Lambillonea*, 100: 601-612.
- PLATIA G. & GUDENZI I. 2002: Revisione dei *Cardiophorus* della Turchia. II. Le specie unicolori (Coleoptera Elateridae). *Boll. Soc. entomol. ital.*, 134 (1): 27-55.
- PLATIA G. & GUDENZI I. 2002: Descrizione di venti nuove specie di Elateridi della regione Paleartica (Insecta Coleoptera Elateridae). *Quaderno di Studi e Notizie di Storia Naturale della Romagna*, 16: 73-98.
- PLATIA G. & GUDENZI I. 2004: Contributo alla conoscenza dei *Dicronychus* Brullé, 1832 della Turcia con descrizione di nuove specie e chiave di determinazione per le specie note (Coleoptera: Elateridae: Cardiophorini). *Bol. S.E.A.*, 34: 9-22.
- PLATIA G. & GUDENZI I. 2005: Description of eleven new species of click-beetles of the Palearctic region, a case of legs teratology and new records of some species of the Italian fauna (Insecta Coleoptera Elateridae). *Quaderno di Studi e Notizie di Storia Naturale della Romagna*, 21: 109-127.
- PLATIA G. & GUDENZI I. 2007: Click-beetles species and records new to the palearctic region (Insecta Coleoptera Elateridae). *Quaderno di Studi e Notizie di Storia Naturale della Romagna*, 24: 79-86.
- PLATIA G., KABALAK M. & SERT O. 2007: Descriptions of two new species of click-beetle (Coleoptera: Elateridae) from Turkey. *Zootaxa*, 1415: 43-47.
- PLATIA G. & KOVANCI B. 2005: A new species of *Athous* Eschscholtz (Coleoptera: Elateridae) from Turkey. *Türkiye entomoloji dergisi*, 29 (1): 5-9.
- PLATIA G. & MARINI M. 1990: Gli Elateridi (Coleoptera) della collezione Andrea Fiori, conservati nel Museo di Zoologia dell'Università di Bologna. *Giornale italiano di Entomologia*, 5: 31-47.
- PLATIA G. & MERTLIK J. 1996: *Anostirus turcicus* sp. n. (Coleoptera: Elateridae) from Turkey. *Klapalekiana*, 32 (3-4): 225-228.
- PLATIA G. & SCHIMMEL R. 1991: Descrizione di nuove specie di Elateridi di Grecia e Turchia con chiave per due gruppi di specie (Coleoptera, Elateridae). *Lavori Soc. Ven. Sc. Nat.*, 16: 131-142.
- PLATIA G. & SCHIMMEL R. 1992: Descrizione di nuove specie di Elateridi di Grecia e Turchia (Coleoptera, Elateridae). *Lavori Soc. Ven. Sc. Nat.*, 17: 87-95.
- PLATIA G. & SCHIMMEL R. 1993: Descrizione di nuove specie di Elateridi di Grecia e Turchia con note su alcuni *Agriotes* Eschscholtz (Coleoptera, Elateridae). *Lavori Soc. Ven. Sc. Nat.*, 18: 51-57.
- PLATIA G. & SCHIMMEL R. 1994: Descrizione di tre nuove specie di Elateridi di Grecia, Turchia e Siria, con note su *Agriotes cibrithorax* Pic, 1910 (Coleoptera, Elateridae). *Lavori Soc. Ven. Sc. Nat.*, 19: 77-82.
- PLATIA G. & TARNAWSKI D. 1998: Description of two new species of click-beetles from Turkey (Coleoptera: Elateridae). *Genus*, 9 (3): 269-274.
- PLATIA G., YILDIRIM E. & KESDEK M. 2007: *Agrypnus sarikamensis* sp. n. and *Athous artvinensis* sp. n., two new species of Elateridae (Coleoptera) from Turkey. *Biocosme Mésogéen*, Nice, 24 (2): 57-63.
- PREISS R. & PLATIA G. 2003: The Click Beetles of Cyprus with descriptions of two new species and notes on species of the genus *Haterumelater* Ôhira, 1968 (Coleoptera: Elateridae). *Z. Arb. Gem. Öst. Ent.*, 55: 97-123.
- REITTER E. 1905: Bestimmungs - Tabelle der europäischen Coleopteren. Elateridae, 1. Theil. Elaterini, Subtribus: Athouina, aus der palaearktischen Fauna. *Vehr. Naturf. Ver. Brünn*, Brünn, 56: 3-122.
- REITTER E. 1910: Beschreibung neuer Coleopteren aus der Familie der Elateriden und Bemerkungen über bekannte Arten. *Wien. Ent. Zeitung*, 29 (5-6): 165-178.
- REITTER E. 1911: Fauna Germanica. Die Käfer Deutschen Reiches. III. Band. *Schr. Dtsch. Lehrerver. Naturk.*, Stuttgart, 26: 436 pp.
- REITTER E. 1918: Bestimmungstabelle der paläarktischen Elater - Arten. *Wien. entomol. Ztg.*, 37 (1-2): 81-105.
- ROUBAL J. 1924: Einige Käferneuheiten. *Entomologischer Anzeiger*, 8: 80-81.
- SAHLBERG J. 1912-1913: Coleoptera mediterranea orientalia, quae in Aegypta, Palaestina, Syria, Caramania atque in Anatolia occidentali anno 1904. *Ofversigt af Finsha Vetenskaps-Societetens Förfärlingar*, 55 (13): 127-132.
- SÁNCHEZ-RUIZ A. & LÖBL I. 2007: Subfamily Cebrioninae, pp. 89-93. In LÖBL I. & SMETANA A. (eds.): *Catalogue of Palaearctic Coleoptera*, Vol. 4. Elateroidea - Derodontoidae - Bostrichoidea - Lymexyloidea - Cleroidea - Cucujoidea. Apollo Books, Stenstrup, 935 pp.
- SCHENKLING S. 1925: *Coleopterorum catalogus, auspiciis et auxilio W. Junk. Pars 80, Elateridae 1*. Berlin, 263 pp.

- SCHENKLING S. 1927: *Coleopterorum catalogus, auspiciis et auxilio W. Junk. Pars 88, Elateridae II.* Berlin, 264-636.
- SCHIMMEL R. 1990: Neue Schnellkäfer, sowie Bemerkungen über bekannte Arten dieser Familie aus der Türkei, aus Spanien und aus Deutschland (Coleoptera, Elateridae). *Entomol. Blätter*, 86 (1-2): 71-84.
- SCHWARZ O. 1992: Revision der paläarktischen Arten der Elateriden - Gattung *Melanotus* Eschsch. *Deutsch. Entomol. Z.*, 1: 145-164.
- SCHWARZ O. 1900: Neue paläarktische Elateriden. *Deutsch. Entomol. Z.*, 1: 98-111.
- SCHWARZ O. 1902: Neue Elateriden. *Stett. Entomol. Ztg.*, 63: 194-316.
- SMETANA A. 2007: Subfamily Lissominae, pp. 207-208. In LÖBL I. & SMETANA A. (eds.): *Catalogue of Palaearctic Coleoptera, Vol. 4. Elateroidea - Derodontoidea - Bosstrochoidea - Lymexyloidea - Cleroidea - Cucujoidae*. Apollo Books, Stenstrup, 935 pp.
- STIBICK J. N. L. 1979: Classification of the Elateridae (Coleoptera). Relationships and classification of the subfamilies and tribes. *Pacific Ins.*, 20 (2/3): 145-186.
- TARNAWSKI D. 1995: A revision of the genus *Selatosomus* Stephens, 1830 (Coleoptera: Elateridae: Athoinae: Ctenicerini). *Genus*, Wroclaw, Supplement: 183 pp.
- TARNAWSKI D. 1996: A world catalogue of Ctenicerini Fleutiaux, 1936 (Coleoptera: Elateridae: Athoinae). *Genus*, Wroclaw, 7 (4): 587-663.
- TARNAWSKI D. 2001: A world catalogue of Ctenicerini Fleutiaux, 1936. Part II (Coleoptera: Elateridae: Athoinae). *Genus*, Wroclaw, 12 (3): 277-323.
- WINKLER A. 1925: *Catalogus Coleopterorum Regionalis Palearcticae*, Wien: 620-663.
- WURST C. 1994: Neue Schnellkäfer der Unterfamilie Athoinae aus dem Taurusgebirge, Südtürkei (Coleoptera: Elateridae). *Mitt. ent. V.*, Stuttgart, 29: 59-66.
- WURST C. 1995: Vier neue Schnellkäfer aus Europa und Kleinasien (Coleoptera: Elateridae). *Entomol. Zeitschrift*, 105 (8): 147-156.
- WURST C. 1995: *Ampedus tingitanus* n. sp. und *Brachygonus hadullanus* n. sp., zwei neue Schnellkäfer aus Nordafrika und Kleinasien (Coleoptera, Elateridae). *Mitt. ent. V.*, Stuttgart, 30: 30-34.
- WURST C. 1997: *Ampedus minos* sp. n. und *Dima raineri* sp. n., zwei neue Schnellkäfer aus Griechenland, mit Bemerkungen zur Synonymie von *Ampedus ocellatus* (Buysson, 1891) (Coleoptera, Elateridae). *Entomologische Blätter*, 93 (1): 1-7.
- WURST C. & SCHIMMEL R. 1995: *Anostirus (Parastirus) nubilosus* n. sp., ein neuer Schnellkäfer aus dem türkischen Kaukasus (Coleoptera, Elateridae). *Mitt. ent. V.*, Stuttgart, 30: 25-29.
- YÜKSEL H. 1970: Türkiye'de bulunan telkurtları (Elateridae) türleri üzerinde sistematik çalışmalar. *Atatürk Üniversitesi Basimevi*, 65 pp.
- ZEISING M. & BRUNNE G. 2003: Neue Elateriden aus Portugal, Marokko und der Türkei (Coleoptera, Elateridae). *Entomologische Blätter für Biologie und Systematik der Käfer*, 98 (3): 189-198.